

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12Q 1/68</b>	<b>A2</b>	(11) International Publication Number: <b>WO 99/66075</b>  (43) International Publication Date: 23 December 1999 (23.12.99)								
<p>(21) International Application Number: PCT/US99/13426</p> <p>(22) International Filing Date: 14 June 1999 (14.06.99)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>09/098,079</td> <td>15 June 1998 (15.06.98)</td> <td>US</td> </tr> <tr> <td>09/097,889</td> <td>15 June 1998 (15.06.98)</td> <td>US</td> </tr> <tr> <td>09/302,681</td> <td>30 April 1999 (30.04.99)</td> <td>US</td> </tr> </table> <p>(71) Applicant (for all designated States except US): MITOKOR [US/US]; 11494 Sorrento Valley Road, San Diego, CA 92121 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): HERRNSTADT, Corinna [DE/US]; 3628 Ruelle DeVillie, San Diego, CA 92130 (US). GHOSH, Soumitra, S. [US/US]; 12334 Pathos Lane, San Diego, CA 92129 (US). CLEVINGER, William [US/US]; 1515 South Melrose Drive #43, Vista, CA 92083 (US). FAHY, Eoin, D. [US/US]; 7171 Canon Hill Way, San Diego, CA 92126 (US). DAVIS, Robert, E. [US/US]; 13272 Glencliff Way, San Diego, CA 92130 (US).</p> <p>(74) Agents: ROSENMAN, Stephen, J. et al.; Seed and Berry LLP, 6300 Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-7092 (US).</p>	09/098,079	15 June 1998 (15.06.98)	US	09/097,889	15 June 1998 (15.06.98)	US	09/302,681	30 April 1999 (30.04.99)	US	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i></p>
09/098,079	15 June 1998 (15.06.98)	US								
09/097,889	15 June 1998 (15.06.98)	US								
09/302,681	30 April 1999 (30.04.99)	US								
<p>(54) Title: DIAGNOSTIC METHOD BASED ON QUANTIFICATION OF EXTRAMITOCHONDRIAL DNA</p> <p>(57) Abstract</p> <p>Compositions and methods based on quantification of extramitochondrial DNA (exmtDNA) sequences are provided that are useful for detecting the presence of or risk for having a disease associated with altered mitochondrial function, and for identifying agents suitable for treating such diseases. The exmtDNA sequences have strong homology to authentic mitochondrial DNA (mtDNA) sequences.</p>										

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## DIAGNOSTIC METHOD BASED ON QUANTIFICATION OF EXTRAMITOCHONDRIAL DNA

### TECHNICAL FIELD

The present invention relates generally to diseases in which altered  
5 mitochondrial function, such as free radical mediated oxidative injury, leads to tissue  
degeneration and, more specifically, to compositions and methods for detecting  
predisposition to such diseases by quantifying extramitochondrial DNA.

### BACKGROUND OF THE INVENTION

A number of degenerative diseases are thought to be caused by or be  
10 associated with alterations in mitochondrial function. These diseases include  
Alzheimer's Disease, diabetes mellitus, Parkinson's Disease, Huntington's disease,  
dystonia, Leber's hereditary optic neuropathy, schizophrenia, and myodegenerative  
disorders such as "mitochondrial encephalopathy, lactic acidosis, and stroke"  
(MELAS), and "myoclonic epilepsy ragged red fiber syndrome" (MERRF). Other  
15 diseases involving altered metabolism or respiration within cells may also be regarded  
as diseases associated with altered mitochondrial function.

Functional mitochondria contain gene products encoded by  
mitochondrial genes situated in mitochondrial DNA (mtDNA) and by  
extramitochondrial genes not situated in the circular mitochondrial genome. The 16.5  
20 kb mtDNA encodes 22 tRNAs, two ribosomal RNAs (rRNA) and only 13 enzymes of  
the electron transport chain (ETC), the elaborate multi-complex mitochondrial assembly  
where, for example, respiratory oxidative phosphorylation takes place. The  
overwhelming majority of mitochondrial structural and functional proteins are encoded  
by extramitochondrial, and in most cases presumably nuclear, genes. Accordingly,  
25 mitochondrial and extramitochondrial genes may interact directly, or indirectly via gene  
products and their downstream intermediates, including metabolites, catabolites,  
substrates, precursors, cofactors and the like. Alterations in mitochondrial function, for  
example impaired electron transport activity, defective oxidative phosphorylation or

increased free radical production, may therefore arise as the result of defective mtDNA, defective extramitochondrial DNA, defective mitochondrial or extramitochondrial gene products, defective downstream intermediates or a combination of these and other factors.

5 Mitochondria are the subcellular organelles that manufacture bioenergetically essential adenosine triphosphate (ATP) by oxidative phosphorylation. Defective mitochondrial activity, including failure at any step of the ETC, may result in the generation of highly reactive free radicals that have the potential of damaging cells and tissues. These free radicals may include reactive oxygen species (ROS) such as  
10 superoxide, peroxynitrite and hydroxyl radicals, and potentially other reactive species that may be toxic to cells. For example, oxygen free radical induced lipid peroxidation is a well established pathogenetic mechanism in central nervous system (CNS) injury, such as that found in a number of degenerative diseases, and in ischemia (*i.e.*, stroke).

There are at least two deleterious consequences of exposure to reactive  
15 free radicals arising from mitochondrial dysfunction that adversely impact the mitochondria themselves. First, free radical mediated damage may inactivate one or more of the myriad proteins of the ETC. According to generally accepted theories of mitochondrial function, proper ETC respiratory activity requires maintenance of an electrochemical potential in the inner mitochondrial membrane by a coupled  
20 chemiosmotic mechanism. Free radical oxidative activity may dissipate this membrane potential, thereby preventing ATP biosynthesis and halting the production of a vital biochemical energy source. In addition, mitochondrial proteins such as cytochrome c and "apoptosis inducing factor" may leak out of the mitochondria after permeability transition and may induce the genetically programmed cell suicide sequence known as  
25 apoptosis or programmed cell death (PCD).

Second, free radical mediated damage may result in catastrophic mitochondrial collapse that has been termed "transition permeability". For example, rapid mitochondrial permeability transition likely entails changes in the inner mitochondrial transmembrane protein adenylate translocase that results in the formation  
30 of a "pore." In any event, because permeability transition is potentiated by free radical

exposure. it may be more likely to occur in the mitochondria of cells from patients having mitochondria associated diseases that are chronically exposed to such reactive free radicals.

Altered mitochondrial function characteristic of the mitochondria associated diseases may also be related to loss of mitochondrial membrane electrochemical potential by mechanisms other than free radical oxidation, and such transition permeability may result from direct or indirect effects of mitochondrial genes, gene products or related downstream mediator molecules and/or extramitochondrial genes. gene products or related downstream mediators, or from other known or unknown causes.

Diabetes mellitus is a common, degenerative disease affecting 5 to 10 percent of the population in developed countries. The propensity for developing diabetes mellitus is reportedly maternally inherited, suggesting a mitochondrial genetic involvement. (Alcolado, J.C. and Alcolado, R., *Br. Med. J.* 302:1178-1180 (1991); Reny, S.L., *International J. Epidem.* 23:886-890 (1994)). Diabetes is a heterogenous disorder with a strong genetic component; monozygotic twins are highly concordant and there is a high incidence of the disease among first degree relatives of affected individuals.

At the cellular level, the degenerative phenotype that may be characteristic of late onset diabetes mellitus includes indicators of altered mitochondrial respiratory function, for example impaired insulin secretion, decreased ATP synthesis and increased levels of reactive oxygen species. Studies have shown that diabetes mellitus may be preceded by or associated with certain related disorders. For example, it is estimated that forty million individuals in the U.S. suffer from late onset impaired glucose tolerance (IGT). IGT patients fail to respond to glucose with increased insulin secretion. A small percentage of IGT individuals (5-10%) progress to insulin deficient non-insulin dependent diabetes (NIDDM) each year. Some of these individuals further progress to insulin dependent diabetes mellitus (IDDM). These forms of diabetes mellitus, NIDDM and IDDM, are associated with decreased release of insulin by pancreatic beta cells and/or a decreased end-organ response to insulin. Other symptoms

of diabetes mellitus and conditions that precede or are associated with diabetes mellitus include obesity, vascular pathologies, peripheral and sensory neuropathies, blindness and deafness.

Parkinson's disease (PD) is a progressive neurodegenerative disorder associated with altered mitochondrial function and characterized by the loss and/or atrophy of dopamine-containing neurons in the *pars compacta* of the *substantia nigra* of the brain. Like Alzheimer's Disease (AD), PD also afflicts the elderly. It is characterized by bradykinesia (slow movement), rigidity and a resting tremor. Although L-Dopa treatment reduces tremors in most patients for a while, ultimately the tremors become more and more uncontrollable, making it difficult or impossible for patients to even feed themselves or meet their own basic hygiene needs.

It has been shown that the neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) induces parkinsonism in animals and man at least in part through its effects on mitochondria. MPTP is converted to its active metabolite, MPP+, in dopamine neurons; it then becomes concentrated in the mitochondria. The MPP+ then selectively inhibits the mitochondrial enzyme NADH:ubiquinone oxidoreductase ("Complex I"), leading to the increased production of free radicals, reduced production of adenosine triphosphate, and ultimately, the death of affected dopamine neurons.

Mitochondrial Complex I is composed of 40-50 subunits; most are encoded by the nuclear genome and seven by the mitochondrial genome. Since parkinsonism may be induced by exposure to mitochondrial toxins that affect Complex I activity, it appears likely that defects in Complex I proteins may contribute to the pathogenesis of PD by causing a similar biochemical deficiency in Complex I activity. Indeed, defects in mitochondrial Complex I activity have been reported in the blood and brain of PD patients (Parker et al., *Am. J. Neurol.* 26:719-723, 1989).

Alzheimer's disease (AD) is a progressive neurodegenerative disorder that is characterized by loss and/or atrophy of neurons in discrete regions of the brain, and that is accompanied by extracellular deposits of  $\beta$ -amyloid and the intracellular accumulation of neurofibrillary tangles. It is a uniquely human disease, affecting over 13 million people worldwide. It is also a uniquely tragic disease. Many individuals

who have lived normal, productive lives are slowly stricken with AD as they grow older, and the disease gradually robs them of their memory and other mental faculties. Eventually, they cease to recognize family and loved ones, and they often require continuous care until their eventual death.

5           There is evidence that defects in oxidative phosphorylation within the mitochondria are at least a partial cause of sporadic AD. The enzyme cytochrome c oxidase (COX), which makes up part of the mitochondrial electron transport chain (ETC), is present in normal amounts in AD patients; however, the catalytic activity of this enzyme in AD patients and in the brains of AD patients at autopsy has been found  
10   to be abnormally low. This suggests that the COX in AD patients is defective, leading to decreased catalytic activity that in some fashion causes or contributes to the symptoms that are characteristic of AD.

Focal defects in energy metabolism in the mitochondria, with accompanying increases in oxidative stress, may be associated with AD. It is well-  
15   established that energy metabolism is impaired in AD brain (Palmer et al., *Brain Res.* 645:338-42, 1994; Pappolla et al., *Am. J. Pathol.* 140:621-28, 1992; Jeandel et al., *Gerontol.* 35:275, 1989; Balazs et al., *Neurochem. Res.* 19:1131-37, 1994; Mecocci et al., *Ann. Neurol.* 36:747-751, 1994; Gsell et al., *J. Neurochem.* 64:1216-23, 1995). For example, regionally specific deficits in energy metabolism in AD brains have been  
20   reported in a number of positron emission tomography studies (Kuhl, et al., *J. Cereb. Blood Flow Metab.* 7:S406, 1987; Grady, et al., *J. Clin. Exp. Neuropsychol.* 10:576-96, 1988; Haxby et al., *Arch. Neurol.* 47:753-60, 1990; Azari et al., *J. Cereb. Blood Flow Metab.* 13:438-47, 1993). Metabolic defects in the temporoparietal neocortex of AD patients apparently presage cognitive decline by several years. Skin fibroblasts from  
25   AD patients display decreased glucose utilization and increased oxidation of glucose, leading to the formation of glycosylation end products (Yan et al., *Proc. Nat. Acad. Sci. USA* 91:7787-91, 1994). Cortical tissue from postmortem AD brain shows decreased activity of the mitochondrial enzymes pyruvate dehydrogenase (Sheu et al., *Ann. Neurol.* 17:444-49, 1985) and  $\alpha$ -ketoglutarate dehydrogenase (Mastrogiacomo et al., *J.*  
30   *Neurochem.* 6:2007-14, 1994), which are both key enzymes in energy metabolism.

Functional magnetic resonance spectroscopy studies have shown increased levels of inorganic phosphate relative to phosphocreatine in AD brain, suggesting an accumulation of precursors that arises from decreased ATP production by mitochondria (Pettegrew et al., *Neurobiol. of Aging* 15:117-32, 1994; Pettigrew et al., *Neurobiol. of*  
5 *Aging* 16:973-75, 1995). In addition, the levels of pyruvate, but not of glucose or lactate, are reported to be increased in the cerebrospinal fluid of AD patients, consistent with defects in cerebral mitochondrial electron transport chain (ETC) activity (Parnetti et al., *Neurosci. Lett.* 199:231-33, 1995).

Signs of oxidative injury are prominent features of AD pathology and, as  
10 noted above, reactive oxygen species (ROS) are critical mediators of neuronal degeneration. Indeed, studies at autopsy show that markers of protein, DNA and lipid peroxidation are increased in AD brain (Palmer et al., *Brain Res.* 645:338-42, 1994; Pappolla et al., *Am. J. Pathol.* 140:621-28, 1992; Jeandel et al., *Gerontol.* 35:275-82, 1989; Balazs et al., *Arch. Neurol.* 4:864, 1994; Mecocci et al., *Ann. Neurol.* 36:747-51,  
15 1994; Smith et al., *Proc. Nat. Acad. Sci. USA* 88:10540-43, 1991). In hippocampal tissue from AD but not from controls, carbonyl formation indicative of protein oxidation is increased in neuronal cytoplasm, and nuclei of neurons and glia (Smith et al., *Nature* 382:120-21, 1996). Neurofibrillary tangles also appear to be prominent sites of protein oxidation (Schweers et al., *Proc. Nat. Acad. Sci. USA* 92:8463, 1995; Blass et al., *Arch. Neurol.* 4:864, 1990). Under stressed and non-stressed conditions incubation  
20 of cortical tissue from AD brains taken at autopsy demonstrate increased free radical production relative to non-AD controls. In addition, the activities of critical antioxidant enzymes, particularly catalase, are reduced in AD (Gsell et al., *J. Neurochem.* 64:1216-23, 1995), suggesting that the AD brain is vulnerable to increased ROS  
25 production. Thus, oxidative stress may contribute significantly to the pathology of mitochondria associated diseases such as AD, where mitochondrial dysfunction and/or elevated ROS may be present.

One hallmark pathology of AD is the death of selected neuronal populations in discrete regions of the brain. Cell death in AD is presumed to be  
30 apoptotic because signs of programmed cell death (PCD) are seen and indicators of



active gliosis and necrosis are not found. (Smale et al., *Exp. Neurolog.* 133:225-230, 1995; Cotman et al., *Molec. Neurobiol.* 10:19-45, 1995.) The consequences of cell death in AD, neuronal and synaptic loss, are closely associated with the clinical diagnosis of AD and are highly correlated with the degree of dementia in AD (DeKosky et al., *Ann. Neurology* 27:457-464, 1990).

Mitochondrial dysfunction is thought to be critical in the cascade of events leading to apoptosis in various cell types (Kroemer et al., *FASEB J.* 9:1277-87, 1995), and may be a cause of apoptotic cell death in neurons of the AD brain. Altered mitochondrial physiology may be among the earliest events in PCD (Zamzami et al., *J. Exp. Med.* 182:367-77, 1995; Zamzami et al., *J. Exp. Med.* 181:1661-72, 1995) and elevated reactive oxygen species (ROS) levels that result from such altered mitochondrial function may initiate the apoptotic cascade (Ausserer et al., *Mol. Cell. Biol.* 14:5032-42, 1994). In several cell types, including neurons, reduction in the mitochondrial membrane potential ( $\Delta\Psi_m$ ) precedes the nuclear DNA degradation that accompanies apoptosis. In cell-free systems, mitochondrial, but not nuclear, enriched fractions are capable of inducing nuclear apoptosis (Newmeyer et al., *Cell* 70:353-64, 1994). Perturbation of mitochondrial respiratory activity leading to altered cellular metabolic states, such as elevated intracellular ROS, may occur in mitochondria associated diseases and may further induce pathogenetic events via apoptotic mechanisms.

Oxidatively stressed mitochondria may release a pre-formed soluble factor that can induce chromosomal condensation, an event preceding apoptosis (Marchetti et al., *Cancer Res.* 56:2033-38, 1996). In addition, members of the Bcl-2 family of anti-apoptosis gene products are located within the outer mitochondrial membrane (Monaghan et al., *J. Histochem. Cytochem.* 40:1819-25, 1992) and these proteins appear to protect membranes from oxidative stress (Korsmeyer et al., *Biochim. Biophys. Act.* 1271:63, 1995). Localization of Bcl-2 to this membrane appears to be indispensable for modulation of apoptosis (Nguyen et al., *J. Biol. Chem.* 269:16521-24, 1994). Thus, changes in mitochondrial physiology may be important mediators of apoptosis. To the extent that apoptotic cell death is a prominent feature of neuronal loss

in AD. mitochondrial dysfunction may be critical to the progression of this disease and may also be a contributing factor in other mitochondria associated diseases.

Regardless of whether a defect underlying a disease associated with altered mitochondrial function may have mitochondrial or extramitochondrial origins, and regardless of whether a defect underlying altered mitochondrial function has been identified, the present invention provides methods that are useful for determining the risk or presence of diseases associated with such altered mitochondrial function, and for identifying agents that are suitable for treating such diseases. In particular, as is elaborated herein below, the present invention provides compositions and methods for the detection of diseases associated with altered mitochondrial function by quantification of unusual mtDNA-like sequences not found in mitochondria and referred to as extramitochondrial DNA (exmtDNA), and other related advantages.

#### SUMMARY OF THE INVENTION

Briefly stated, the present invention is directed to compositions and methods useful for detecting mitochondria associated diseases and involving extramitochondrial DNA (exmtDNA) sequences that are highly homologous to mitochondrial DNA (mtDNA). In one aspect the invention provides a method for determining the risk for or presence of a disease associated with altered mitochondrial function in a first subject suspected of having or being at risk for having such a disease, by comparing a ratio  $r$  for each of a first and a second biological sample containing extramitochondrial DNA and mitochondrial DNA, the first biological sample being obtained from the first subject and the second sample being obtained from a second subject known to be free of a risk or presence of a disease associated with altered mitochondrial function, using the formula:

$$r = x/(x + y)$$

wherein  $x$  is the amount of extramitochondrial DNA in a sample, and  $y$  is the amount of mitochondrial DNA in the sample; and therefrom determining the risk or presence of the disease. In an embodiment of the invention, the ratio  $r$  is calculated by a method

that comprises contacting a biological sample containing extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in the extramitochondrial DNA and present in the mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of  
5 the primer to the extramitochondrial DNA and to the mitochondrial DNA; and detecting hybridization of the primer to the extramitochondrial DNA and to the mitochondrial DNA, in order to therefrom quantify the extramitochondrial DNA and the mitochondrial DNA.

In another embodiment, the ratio  $r$  is calculated by a method comprising  
10 contacting a sample containing amplified extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in the amplified extramitochondrial DNA and present in the amplified mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of the primer to the extramitochondrial DNA and to the  
15 mitochondrial DNA; and detecting hybridization of the primer to the extramitochondrial DNA and to the mitochondrial DNA, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA. In another embodiment of this aspect of the invention the ratio  $r$  is calculated by a method comprising contacting a biological sample containing extramitochondrial DNA and mitochondrial DNA with an  
20 oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in the extramitochondrial DNA and present in the mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of said primer to the extramitochondrial DNA and to the mitochondrial DNA; and detecting hybridization and extension of the primer to the extramitochondrial DNA to produce a  
25 first product and hybridization and extension of the primer to the mitochondrial DNA to produce a second product distinguishable from the first product, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA.

In another embodiment of this aspect of the invention the ratio  $r$  is calculated by a method comprising contacting a sample containing amplified  
30 extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer

having a nucleotide sequence that is complementary to a sequence present in the amplified extramitochondrial DNA and present in the amplified mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of the primer to the extramitochondrial DNA and to the mitochondrial DNA; and detecting hybridization  
5 and extension of the primer to the extramitochondrial DNA to produce a first product and hybridization and extension of the primer to the mitochondrial DNA to produce a second product distinguishable from the first product, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA.

In another embodiment of this aspect of the invention the biological  
10 sample is treated by heating it in water to lyse cells contained in the sample, and then extracting cellular DNA from the lysed cells using an aqueous DNA extraction procedure. In certain embodiments of the invention the sample comprises a crude buffy coat fraction of whole blood. In certain other embodiments of the invention, the method further comprises the step of determining the ApoE genotype of the first subject and  
15 correlating said genotype with the risk or presence of disease. In some embodiments of the invention, the disease associated with altered mitochondrial function may be Alzheimer's Disease, Huntington's Disease, Parkinson's Disease, dystonia, schizophrenia, non-insulin dependent diabetes mellitus, mitochondrial encephalopathy, lactic acidosis, and stroke, myoclonic epilepsy ragged red fiber syndrome, and Leber's  
20 hereditary optic neuropathy.

Another aspect of the invention provides a method for quantifying extramitochondrial DNA, comprising: contacting a sample containing extramitochondrial DNA with an oligonucleotide primer having a nucleotide sequence complementary to at least a portion of the extramitochondrial DNA under conditions  
25 and for a time sufficient to allow hybridization of the primer to the extramitochondrial DNA; and detecting hybridization of the primer to the extramitochondrial DNA, and therefrom quantifying the extramitochondrial DNA.

It is another aspect of the invention to provide a method for quantifying extramitochondrial DNA, comprising: contacting a sample containing  
30 extramitochondrial DNA with an oligonucleotide primer having a nucleotide sequence

complementary to at least a portion of the extramitochondrial DNA under conditions and for a time sufficient to allow hybridization of the primer to the extramitochondrial DNA; and detecting hybridization and extension of the primer to the extramitochondrial DNA to produce a product, and therefrom quantifying the extramitochondrial DNA.

5 Another aspect of the invention provides a method for quantifying extramitochondrial DNA, comprising: contacting a sample containing amplified extramitochondrial DNA with an oligonucleotide primer having a nucleotide sequence complementary to at least a portion of the extramitochondrial DNA under conditions and for a time sufficient to allow hybridization of the primer to the extramitochondrial  
10 DNA; and detecting hybridization of the primer to the extramitochondrial DNA, therefrom quantifying the extramitochondrial DNA.

In yet another aspect of the invention, a method is provided for quantifying extramitochondrial DNA by contacting a sample containing amplified extramitochondrial DNA with an oligonucleotide primer having a nucleotide sequence  
15 complementary to at least a portion of the extramitochondrial DNA under conditions and for a time sufficient to allow hybridization of the primer to the extramitochondrial DNA; and detecting hybridization and extension of the primer to the extramitochondrial DNA to produce a product, and therefrom quantifying the extramitochondrial DNA.

In one embodiment the extramitochondrial DNA is amplified by  
20 polymerase chain reaction, transcriptional amplification systems or self-sustained sequence replication. In certain embodiments of the various aspects of the invention, a single oligonucleotide primer is used. In certain embodiments of the invention a primer extension assay is used. In certain embodiments of the invention, the step of detecting may be by polymerase chain reaction, primer extension assay, ligase chain reaction or  
25 restriction fragment length polymorphism analysis.

In certain embodiments of the invention, the amount of extramitochondrial DNA in a biological sample is quantified by determining the presence in the sample of a nucleotide sequence that may be SEQ ID NO:1, a portion of SEQ ID NO:1, SEQ ID NO:3, a portion of SEQ ID NO:3, an extramitochondrial DNA  
30 sequence comprising a nucleic acid sequence that (i) corresponds to at least a portion of

SEQ ID NO:2 and (ii) contains at least one nucleotide substitution of Figure 4 at a corresponding nucleotide position, or an extramitochondrial DNA sequence comprising a nucleic acid sequence that (i) corresponds to at least a portion of SEQ ID NO:2 and (ii) contains at least one nucleotide deletion of Figure 4 at a corresponding nucleotide position. In one embodiment the portion of the nucleotide sequence of SEQ ID NO:1 corresponds to a portion of the nucleotide sequence of SEQ ID NO:2 encoding a mitochondrial cytochrome c oxidase. In another embodiment the portion of SEQ ID NO:1 corresponds to a portion of a mitochondrial cytochrome c oxidase encoding sequence that may be portion of a cytochrome c oxidase 1 (CO1) encoding sequence or a portion of a cytochrome c oxidase 2 (CO2) encoding sequence. In still other embodiments, the portion of the nucleotide sequence of SEQ ID NO:1 corresponds to a portion of the nucleotide sequence of SEQ ID NO:2 encoding a mitochondrial ATP synthetase subunit. In other embodiments, the portion of SEQ ID NO:1 corresponds to a portion of a mitochondrial ATP synthetase subunit encoding sequence that may be a portion of a sequence encoding ATP synthetase subunit 6 or a portion of a sequence encoding ATP synthetase subunit 8.

In some embodiments the nucleotide sequence of SEQ ID NO:1 corresponds to a portion of SEQ ID NO:2 that may be a portion of a sequence encoding ND1, a sequence encoding a portion of ND2 or a sequence encoding a portion of CO3. In other embodiments, the portion of the nucleotide sequence of SEQ ID NO:3 corresponds to a portion of the nucleotide sequence of SEQ ID NO:2 encoding a mitochondrial ATP synthetase subunit, which in some embodiments may further be a portion of a sequence encoding ATP synthetase subunit 6 or a portion of a sequence encoding ATP synthetase subunit 8. In still other embodiments, the nucleotide sequence of SEQ ID NO:1 corresponds to a portion of the nucleotide sequence of SEQ ID NO:2 encoding a mitochondrial tRNA, while in yet other embodiments the portion of the nucleotide sequence of SEQ ID NO:3 corresponds to a portion of the nucleotide sequence of SEQ ID NO:2 encoding a mitochondrial tRNA.

In another aspect the invention provides an isolated nucleic acid comprising all or a portion of the nucleotide sequence of SEQ ID NO:1 or a

complementary sequence thereto. In another aspect the invention provides an isolated nucleic acid comprising all or a portion of a nucleotide sequence of SEQ ID NO:1 or a complementary sequence thereto, wherein the sequence of the isolated nucleic acid differs by at least one nucleotide from the corresponding sequence of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:2 or a complementary sequence thereto. In another aspect the invention provides an isolated nucleic acid comprising all or a portion of the nucleotide sequence of SEQ ID NO:3 or a complementary sequence thereto. In another aspect the invention provides an isolated nucleic acid comprising all or a portion of a nucleotide sequence of SEQ ID NO:3 or a complementary sequence thereto, wherein the sequence of the isolated nucleic acid differs by at least one nucleotide from the corresponding sequence of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:2 or a complementary sequence thereto.

In another aspect the invention provides a method for determining the risk or presence of a disease associated with altered mitochondrial function in a subject suspected of having or being at risk for having such a disease, by quantifying the amount of extramitochondrial DNA and the amount of mitochondrial DNA in a biological sample from the subject, and therefrom determining the risk or presence of the disease. It is another aspect of the invention to provide a method for determining the risk or presence of a disease associated with altered mitochondrial function in a first subject suspected of having or being at risk for having such a disease, by comparing the amount of extramitochondrial DNA and the amount of mitochondrial DNA in a biological sample from the first subject to the amount of extramitochondrial DNA and the amount of mitochondrial DNA in a biological sample from a second subject, and therefrom determining the risk or presence of the disease. In another aspect the invention provides a method for determining the risk or presence of a disease associated with altered mitochondrial function in a first subject suspected of having or being at risk for having such a disease, by quantifying the amount of extramitochondrial DNA and the amount of mitochondrial DNA in a biological sample from the subject and comparing the amount of extramitochondrial DNA and the amount of mitochondrial DNA to the amount of extramitochondrial DNA and the amount of mitochondrial DNA

in a biological sample from a second subject known to be free of a risk or presence of a disease associated with altered mitochondrial function. and therefrom determining the risk or presence of the disease.

Another aspect of the invention provides a method of monitoring a  
5 telomere by detecting in a telomeric region the presence of a nucleic acid molecule comprising all or a portion of SEQ ID NO:1 or a complementary portion thereto. In one embodiment, the detected nucleic acid molecule comprises an exmtDNA sequence. In another aspect, the invention provides a method of monitoring a telomere by detecting in a telomeric region the presence of a nucleic acid molecule comprising all or a portion  
10 of SEQ ID NO:3 or a complementary portion thereto.

Turning to another aspect, the invention provides a method of identifying  
an agent suitable for treating a disease associated with altered mitochondrial function,  
by comparing a ratio  $r$  from a sample obtained before contacting a biological source with a candidate agent to the ratio  $r$  from a sample obtained after contacting the  
15 biological source with the candidate agent, said ratio  $r$  calculated using the formula:

$$r = x/(x + y)$$

wherein  $x$  is the amount of extramitochondrial DNA in a sample, and  $y$  is the amount of  
20 mitochondrial DNA in the sample; and therefrom determining the suitability of said candidate agent for treating a disease associated with altered mitochondrial function. In one embodiment, the biological sample may be a crude buffy coat fraction of whole blood. In another embodiment, the biological sample is treated by heating in water to lyse cells contained in the sample, and then extracting cellular DNA from lysed cells  
25 using an aqueous DNA extraction procedure. In another embodiment, the ratio  $r$  is calculated by contacting a sample containing extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in the extramitochondrial DNA and present in the mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of  
30 the primer to the extramitochondrial DNA and to the mitochondrial DNA; and detecting hybridization of the primer to the extramitochondrial DNA and to the mitochondrial



DNA, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA to calculate the ratio  $r$ .

In another embodiment of the invention, the ratio  $r$  is calculated by contacting a sample containing extramitochondrial DNA and mitochondrial DNA with  
5 an oligonucleotide primer having a nucleotide sequence that is complementary to a  
sequence present in the extramitochondrial DNA and present in the mitochondrial  
DNA, under conditions and for a time sufficient to allow hybridization of the primer to  
the extramitochondrial DNA and to the mitochondrial DNA; and detecting  
hybridization and extension of the primer to the extramitochondrial DNA to produce a  
10 first product and hybridization and extension of the primer to the mitochondrial DNA to  
produce a second product distinguishable from the first product, and therefrom  
quantifying the extramitochondrial DNA and the mitochondrial DNA to calculate the  
ratio. In another embodiment, the ratio  $r$  is calculated by contacting a sample  
containing amplified extramitochondrial DNA and mitochondrial DNA with an  
15 oligonucleotide primer having a nucleotide sequence that is complementary to a  
sequence present in said amplified extramitochondrial DNA and present in said  
amplified mitochondrial DNA, under conditions and for a time sufficient to allow  
hybridization of said primer to the extramitochondrial DNA and to the mitochondrial  
DNA; and detecting hybridization of the primer to the extramitochondrial DNA and to  
20 the mitochondrial DNA, and therefrom quantifying the extramitochondrial DNA and  
the mitochondrial DNA to calculate the ratio  $r$ .

In yet another embodiment, the ratio  $r$  is calculated by contacting a  
sample containing amplified extramitochondrial DNA and mitochondrial DNA with an  
oligonucleotide primer having a nucleotide sequence that is complementary to a  
25 sequence present in the amplified extramitochondrial DNA and present in the amplified  
mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of  
said primer to the extramitochondrial DNA and to the mitochondrial DNA; and  
detecting hybridization and extension of the primer to the extramitochondrial DNA to  
produce a first product and hybridization and extension of the primer to the  
30 mitochondrial DNA to produce a second product distinguishable from said first product,

and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA to calculate the ratio.

In another embodiment of the invention, comparing the ratio  $r$  from a sample obtained before contacting a biological source with a candidate agent to the ratio  $r$  from a sample obtained after contacting the biological source with the candidate agent comprises determination of the presence in the sample of a nucleotide sequence of SEQ ID NO:1 or portion thereof, or a nucleotide sequence of SEQ ID NO:3 or a portion thereof, or an extramitochondrial DNA sequence comprising a nucleic acid sequence that (i) corresponds to at least a portion of SEQ ID NO:2 and (ii) contains at least one nucleotide substitution of Figure 4 at a corresponding nucleotide position, or an extramitochondrial DNA sequence comprising a nucleic acid sequence that (i) corresponds to at least a portion of SEQ ID NO:2 and (ii) contains at least one nucleotide deletion of Figure 4 at a corresponding nucleotide position. In another embodiment, the nucleotide sequence of SEQ ID NO:1 or a portion thereof corresponds to a mitochondrial cytochrome c oxidase encoding sequence of SEQ ID NO:2 or a portion thereof. In another embodiment, the mitochondrial cytochrome c oxidase encoding sequence of SEQ ID NO:2 or a portion thereof is a sequence encoding CO I or a portion thereof, or a sequence encoding CO2 or a portion thereof. In another embodiment, the nucleotide sequence of SEQ ID NO:1 or portion thereof, or the nucleotide sequence of SEQ ID NO:3 or portion thereof corresponds to a mitochondrial ATP synthetase subunit encoding sequence of SEQ ID NO:2 or a portion thereof. In another embodiment, the mitochondrial ATP synthetase subunit encoding sequence of SEQ ID NO:2 or a portion thereof may be a sequence encoding ATP synthetase subunit 6 or a portion thereof, or a sequence encoding ATP synthetase subunit 8 or a portion thereof. In another embodiment, the nucleotide sequence of SEQ ID NO:1 corresponds to a sequence of SEQ ID NO:2 or a portion thereof that may be a sequence encoding a truncated NADH dehydrogenase subunit 1 or a portion thereof, a sequence encoding NADH dehydrogenase subunit 2 or a portion thereof or a sequence encoding truncated CO3 or a portion thereof.

In other embodiments of the invention, the disease associated with altered mitochondrial function may be Alzheimer's Disease, Huntington's Disease, Parkinson's Disease, dystonia, schizophrenia, non-insulin dependent diabetes mellitus, mitochondrial encephalopathy, lactic acidosis, and stroke, myoclonic epilepsy ragged  
5 red fiber syndrome, or Leber's hereditary optic neuropathy.

In another aspect, the invention provides a method of identifying an agent suitable for treating a subject suspected of being at risk for having a disease associated with altered mitochondrial function, by determining the apolipoprotein E genotype of the subject; comparing a ratio  $r$  in a biological sample obtained from the  
10 subject before contacting the sample with a candidate agent to the ratio  $r$  in a biological sample obtained from the subject after contacting the sample with a candidate agent, the ratio  $r$  calculated using the formula:

$$r = x / (x + y)$$

15

wherein  $x$  is the amount of extramitochondrial DNA in the sample, and  $y$  is the amount of mitochondrial DNA in the sample; and therefrom determining the suitability of said candidate agent for treating the disease associated with altered mitochondrial function. In another embodiment, the disease associated with altered mitochondrial function is  
20 Alzheimer's disease.

It is another aspect of the invention to provide a method of correlating a ratio  $r$  with the suitability of an agent for treating Alzheimer's disease in a subject, by determining a ratio  $r$  in a biological sample obtained from the subject, said ratio  $r$  calculated using the formula:

25

$$r = x / (x + y)$$

wherein  $x$  is the amount of extramitochondrial DNA in the sample, and  $y$  is the amount of mitochondrial DNA in the sample; contacting said subject with a candidate agent and  
30 evaluating the subject for alterations in the AD disease state, and therefrom correlating the suitability of the agent for treating AD in the subject with  $r$ . In another embodiment, the apolipoprotein E genotype of the subject is determined, and therefrom

the suitability of the agent for treating AD in the subject is correlated with  $r$  and with the apolipoprotein E genotype.

These and other aspects of the present invention will become evident upon reference to the following detailed description and attached drawings. In addition, various references are set forth herein which describe in more detail certain aspects of this invention, and are therefore incorporated by reference in their entirety.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the nucleotide sequence of SEQ ID NO:1.

Figure 2 depicts the nucleotide sequence of SEQ ID NO:2.

10 Figure 3 depicts the nucleotide sequence of SEQ ID NO:3.

Figure 4 depicts human extramitochondrial DNA nucleotide substitutions and deletions relative to the corresponding human mtDNA sequence of SEQ ID NO:2.

15 Figure 5 shows radiation hybrid chromosomal mapping of an extramitochondrial DNA sequence.

Figure 6 depicts detection of mtDNA and exmtDNA by Southern blot hybridization.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed generally to compositions and methods for diagnosing the risk or presence of a disease associated with altered mitochondrial function in a subject, and to compositions and methods for the identification of agents that may be suitable for treating a disease associated with altered mitochondrial function. The invention utilizes a ratio,  $r$ , that may be useful for pharmacogenomic purposes, for example to stratify patient populations according to the suitability of particular therapeutic agents for use in such populations. The ratio  $r$  is the ratio of the amount of exmtDNA in a biological sample relative to the sum of the amount of exmtDNA plus mtDNA in the sample. As expressed quantitatively, the ratio  $r$  may be calculated using the formula:

$$r = x/(x + y)$$

wherein

$x$  is the amount of exmtDNA in a sample. and

5  $y$  is the amount of mtDNA in the sample.

In various aspects of the invention, as elaborated more fully herein, quantification of  $x$  and  $y$  provide, through calculation of  $r$ , parameters useful in diagnosis of a disease associated with altered mitochondrial function and in screening assays for agents that may be suitable for the treatment of such a disease.

10 As discussed above, "altered mitochondrial function" may refer to any condition or state, including those that accompany a disease, where any structure or activity that is directly or indirectly related to a mitochondrial function has been changed. Altered mitochondrial function may have its origin in extramitochondrial structures or events as well as in mitochondrial structures or events, in direct  
15 interactions between mitochondrial and extramitochondrial genes and/or their gene products, or in structural or functional changes that occur as the result of interactions between intermediates that may be formed as the result of such interactions, including metabolites, catabolites, substrates, precursors, cofactors and the like.

Also as discussed above, altered mitochondrial function may include  
20 (but need not be limited to) altered respiratory or metabolic activity in some or all cells of a biological source. For example, markedly impaired ETC activity may be an example of altered mitochondrial function, as may be generation of increased ROS or defective oxidative phosphorylation. As further examples, altered mitochondrial membrane potential, induction of apoptotic pathways and formation of atypical  
25 chemical and biochemical crosslinked species within a cell, whether by enzymatic or non-enzymatic mechanisms, may all be regarded as indicative of altered mitochondrial function. Without wishing to be bound by theory, alterations in the ratio  $r$  are believed to reflect chemical changes within affected cells that quantitatively influence recoveries of exmtDNA and/or mtDNA. For example, biochemical crosslinking events may result  
30 in the formation of DNA aggregates, DNA adducts or other molecular species that affect DNA recoveries following extraction procedures.

According to the present invention, alterations in the ratio  $r$  as defined above provide a novel and useful parameter for diagnosing the risk or presence of a disease associated with altered mitochondrial function in a subject, and for identifying agents that may be suitable for treating a disease associated with altered mitochondrial function. As discussed above, a number of diseases, including several degenerative diseases, are associated with alterations in mitochondrial function. Further, detection of an appropriate parameter of altered mitochondrial function can provide preclinical evidence for a risk of or predisposition to a disease.

Determination of the ratio  $r$  involves quantification of exmtDNA ( $x$ ) and mtDNA ( $y$ ) that may be based on strong but not necessarily absolute nucleotide sequence conservation when corresponding portions of mtDNA and exmtDNA are compared, as discussed herein. In most preferred embodiments of the invention, determination of  $r$  is accomplished by detecting minor nucleotide sequence differences in highly conserved mtDNA and exmtDNA regions, as elaborated below. The invention provides compositions and methods that include the use of nucleic acid molecules, or portions thereof, having nucleotide sequences that are found in the human mtDNA sequence SEQ ID NO:2 (Anderson et al., *Nature* 290:457, 1981) and fragments of SEQ ID NO:2 that are suitable for use as oligonucleotide primers in nucleic acid primer extension or amplification techniques, as hybridization probes for the detection of complementary nucleotide sequences in a sample or for any number of additional uses that are well known to those familiar with the art. ExmtDNA may be nuclear DNA, including chromosomal and non-chromosomal DNA, or non-nuclear extramitochondrial DNA that may be from any subcellular compartment, provided it is not mtDNA.

Nucleic acid sequences within the scope of the invention include isolated DNA and RNA sequences that specifically hybridize under conditions of moderate or high stringency to exmtDNA nucleotide sequences, including exmtDNA sequences disclosed herein or fragments thereof, and their complements. As used herein, conditions of moderate stringency, as known to those having ordinary skill in the art, and as defined by Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed.

Vol. 1, pp. 1.101-104. Cold Spring Harbor Laboratory Press (1989). include use of a prewashing solution for the nitrocellulose filters 5X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0), hybridization conditions of 50% formamide, 6X SSC at 42°C (or other similar hybridization solution). and washing conditions of about 50-60°C, 0.5X SSC, 0.1% SDS. Conditions of high stringency are defined as hybridization conditions as above, and with washing at 60-68°C, 0.2X SSC, 0.1% SDS. In other embodiments, hybridization to an exmtDNA nucleotide sequence may be at normal stringency, which is approximately 25-30°C below T<sub>m</sub> of the native duplex (e.g., 5X SSPE, 0.5% SDS, 5X Denhardt's solution, 50% formamide, at 42°C or equivalent conditions), at low stringency hybridizations, which utilize conditions approximately 40°C below T<sub>m</sub>, or at high stringency hybridizations, which utilize conditions approximately 10°C below T<sub>m</sub>. The skilled artisan will recognize that the temperature, salt concentration, and chaotrope composition of hybridization and wash solutions may be adjusted as necessary according to factors such as the length and nucleotide base composition of the probe. (See also, e.g., Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing, 1987.)

An "isolated nucleic acid molecule" refers to a polynucleotide molecule in the form of a separate fragment or as a component of a larger nucleic acid construct, that has been separated from its source cell (including the chromosome it normally resides in) at least once, preferably in a substantially pure form. Isolated nucleic acids may be nucleic acids having particular disclosed nucleotide sequences or may be regions, portions or fragments thereof. Those having ordinary skill in the art are able to prepare isolated nucleic acids having the complete nucleotide sequence, or the sequence of any portion of a particular isolated nucleic acid molecule, when provided with the appropriate nucleic acid sequence information as disclosed herein. Nucleic acid molecules may be comprised of a wide variety of nucleotides, including DNA, RNA, nucleotide analogues such as phosphorothioates or peptide nucleic acids, or other analogues with which those skilled in the art will be familiar, or some combination of these.

The present invention, as described herein, provides exmtDNA sequences and isolated exmtDNA nucleic acid molecules. exmtDNA may be isolated from genomic DNA, typically by first generating an appropriate DNA library through techniques for constructing libraries that are known in the art (*see* Sambrook et al., 5 *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, 1989) or purchased from commercial sources (*e.g.*, Clontech, Palo Alto, California). Briefly, genomic DNA libraries can be constructed in chromosomal vectors, such as YACs (yeast artificial chromosomes), bacteriophage vectors, such as pBeloBAC11,  $\lambda$ EMBL3,  $\lambda$ gt10, cosmids, or plasmids. Alternatively, isolated exmtDNA may be prepared by 10 preferentially amplifying exmtDNA sequences present in biological samples using, for example, DNA amplification methodologies such as PCR or other amplification techniques that are well known in the art, with suitable oligonucleotide primers complementary to exmtDNA sequences as disclosed herein.

In one embodiment, known mtDNA sequences derived from SEQ ID 15 NO:2 (Anderson et al., *Nature* 290:457, 1981) may be utilized to design oligonucleotide hybridization probes suitable for screening genomic libraries. Preferably, such oligonucleotide probes are 18-30 bases in length and have sequences that, under the hybridization conditions selected, hybridize to complementary exmtDNA sequences lacking nucleotide substitutions, insertions or deletions ("mutations") relative to the 20 corresponding region of the mtDNA sequence of SEQ ID NO:2.

Portions of an exmtDNA sequence and the mtDNA sequence of SEQ ID NO:2 are regarded as "corresponding" nucleic acid sequences, regions, fragments or the like, based on the convention for numbering mtDNA nucleic acid positions according to SEQ ID NO:2 (Anderson et al., *Nature* 290:457, 1981), wherein an exmtDNA sequence 25 is aligned with the mtDNA sequence of SEQ ID NO:2 such that at least 70%, preferably at least 80% and more preferably at least 90% of the nucleotides in a given sequence of at least 20 consecutive nucleotides of a sequence are identical. In certain preferred embodiments, an exmtDNA sequence is greater than 95% identical to a corresponding mtDNA sequence. In certain particularly preferred embodiments, an exmtDNA 30 sequence is identical to a corresponding mtDNA sequence. Those oligonucleotide



probes having sequences that are identical in corresponding regions of mtDNA and exmtDNA may be identified and selected following hybridization target DNA sequence analysis, to verify the absence of mutations in the target exmtDNA sequence relative to the primer mtDNA-derived sequence.

5 To facilitate hybridization detection, the oligonucleotide may be conveniently labeled, generally at the 5' end, with a reporter molecule, such as a radionuclide, *e.g.*, <sup>32</sup>P, enzymatic label, protein label, fluorescent label, biotin or other suitable labeling moieties known in the art. Such libraries are then generally plated as phage or colonies, depending upon the vector used. Subsequently, a plate replica to  
10 which the colonies or phage have been transferred, such as a nitrocellulose or nylon membrane or the like, is probed to identify candidate clones that contain the exmtDNA sequence. Such candidates may be verified as containing exmtDNA by any of various means including, for example, DNA sequence analysis or hybridization with a second, non-overlapping probe selected as described above to hybridize with target exmtDNA  
15 sequences lacking nucleotide substitutions, deletions or insertions relative to the corresponding portion of the mtDNA sequence of SEQ ID NO:2.

Once a library is identified as containing exmtDNA, the exmtDNA can be isolated by amplification. Briefly, when using genomic library DNA as a template, amplification primers are designed based upon known mtDNA sequences (SEQ ID  
20 NO:2) and primer "walking" is used to select primers that anneal to exmtDNA regions that are identical to mtDNA sequences. The primers preferably have a GC content of about 50% and contain restriction sites to facilitate cloning. Primers do not have self-complementary sequences, nor do they contain complementary sequences at their 3' end (to prevent primer-dimer formation). The primers are annealed to genomic DNA and  
25 sufficient amplification cycles are performed to yield a product readily visualized by gel electrophoresis and staining. The amplified fragment is purified and inserted into a vector, such as  $\lambda$ gt10 or pBS(M13+), and propagated. Confirmation of the nature of the fragment is obtained by DNA sequence analysis.

As an example of detection of mtDNA-like sequences in a DNA library,  
30 an oligonucleotide having a nucleotide sequence present in a portion of any human

mtDNA gene, preferably one of the human mtDNA encoded genes NADH dehydrogenase subunit 1 (ND1), NADH dehydrogenase subunit 2 (ND2), NADH dehydrogenase subunit 3 (ND3), NADH dehydrogenase subunit 4 (ND4), NADH dehydrogenase subunit 4L (ND4L), NADH dehydrogenase subunit 5 (ND5), NADH dehydrogenase subunit 6 (ND6), tRNA<sup>Lys</sup>, tRNA<sup>Ile</sup>, tRNA<sup>Gln</sup>, tRNA<sup>Met</sup>, tRNA<sup>Cys</sup>, tRNA<sup>Tyr</sup>, tRNA<sup>Asp</sup> or cytochrome c oxidase 3 (CO3) and more preferably one of the human mtDNA encoded genes CO1, CO2, ATPase 8 or ATPase 6, may be labeled and used as a probe on a human genomic DNA library. An initial hybridization at normal stringency may yield candidate clones or fragments. If no hybridization is initially observed, varying degrees of stringency may be used. (See Sambrook et al., Ausubel et al., *supra*, and other well-known sources for stringency conditions.)

Where it is advantageous to use oligonucleotide primers according to the present invention, such primers may be 10-60 nucleotides in length, preferably 15-35 nucleotides and still more preferably 18-30 nucleotides in length. Primers as described above for use in isolating exmtDNA from genomic DNA may also be useful in the present invention for quantifying mtDNA and exmtDNA by any of a variety of techniques well known in the art for determining the amount of specific nucleic acid target sequences present in a sample based on specific hybridization of a primer to the target sequence. Optionally, in certain of these techniques, hybridization precedes nucleotide polymerase catalyzed extension of the primer using the strand containing the target sequence as a template, and/or ligation of oligonucleotides hybridized to adjacent target sequences, and embodiments of the invention using primer extension are particularly preferred. For examples of references on such quantitative detection techniques, including those that may be used to detect nucleotide insertions, substitutions or deletions in a portion of an exmtDNA sequence site near an oligonucleotide primer target hybridization site that corresponds to a portion of the mtDNA sequence of SEQ ID NO:2, and further including those that involve primer extension, see U.S. 5,760,205 and the references cited therein, all of which are hereby incorporated by reference, and see also, for example, Botstein et al. (*Am. J. Hum. Gen.* 32:314, 1980), Gibbs et al. (*Nucl. Ac. Res.* 17:2437, 1989), Newton et al. (*Nucl. Ac. Res.*

17:2503, 1989), Grossman et al. (*Nucl. Ac. Res.* 22:4527, 1994), and Saiki et al. (*Proc. Nat. Acad. Sci.* 86:6230, 1989), all of which are hereby incorporated by reference. A particularly useful method for this purpose is the primer extension assay disclosed by Fahy et al. (*Nucl. Acids Res.* 25:3102, 1997) and by Ghosh et al. (*Am. J. Hum. Genet.* 58:325, 1996), both of which references are hereby incorporated in their entireties, as is Krok et al. (*Hum. Molec. Genet.* 1:391, 1995) which teaches modification of primer extension reactions to detect multiple nucleotide substitutions, insertions, deletions or other mutations. Other examples of useful techniques for quantifying the presence of specific nucleic acid target sequences in a sample include but need not be limited to labeled probe hybridization to the target nucleic acid sequences with or without first partially separating target nucleic acids from other nucleic acids present in the sample.


Examples of other useful techniques for determining the amount of specific nucleic acid target sequences present in a sample based on specific hybridization of a primer to the target sequence include specific amplification of target nucleic acid sequences and quantification of amplification products, including but not limited to polymerase chain reaction (PCR, Gibbs et al., *Nucl. Ac. Res.* 17:2437, 1989), transcriptional amplification systems, strand displacement amplification and self-sustained sequence replication (3SR, Ghosh et al, in *Molecular Methods for Virus Detection*, 1995 Academic Press, NY, pp. 287-314), the cited references for which are hereby incorporated in their entireties. Examples of other useful techniques include ligase chain reaction, single stranded conformational polymorphism analysis, Q-beta replicase assay, restriction fragment length polymorphism (RFLP, Botstein et al., *Am. J. Hum. Gen.* 32:314, 1980) analysis and cycled probe technology, as well as other suitable methods that will be known to those familiar with the art.

In a particularly preferred embodiment of the invention, primer extension is used to quantify exmtDNA and mtDNA present in a biological sample. (Ghosh et al., *Am. J. Hum. Genet.* 58:325, 1996) This embodiment may offer certain advantages by permitting both exmtDNA and mtDNA to be simultaneously quantified using a single oligonucleotide primer capable of hybridizing to a complementary nucleic acid target sequence that is present in a defined region of mtDNA and in a corresponding region of

a exmtDNA sequence. Without wishing to be bound by theory, the use of a single primer for quantification of exmtDNA and of mtDNA is believed to avoid uncertainties associated with potential disparities in the relative hybridization properties of multiple primers and may offer other advantages. Where such a target sequence is situated adjacent to an exmtDNA nucleotide sequence position that is a nucleotide substitution, insertion or deletion relative to the corresponding mtDNA sequence position, primer extension assays may be designed such that oligonucleotide extension products of primers hybridizing to mtDNA are of different lengths than oligonucleotide extension products of primers hybridizing to exmtDNA. Accordingly, the amount of exmtDNA in a sample and the amount of mtDNA in the sample may be determined by quantification of distinct extension products that are separable on the basis of sequence length or molecular mass, for purposes of calculating the ratio  $r$  as described above.

Sequence length or molecular mass of primer extension assay products may be determined using any known method for characterizing the size of nucleic acid sequences with which those skilled in the art are familiar. In a preferred embodiment, primer extension products are characterized by gel electrophoresis. In another preferred embodiment, primer extension products are characterized by mass spectrometry (MS), which may further include matrix assisted laser desorption ionization/ time of flight (MALDI-TOF) analysis or other MS techniques known to those having skill in the art. See, for example, U.S. 5,622,824, U.S. 5,605,798 and U.S. 5,547,835, all of which are hereby incorporated by reference in their entireties. In another preferred embodiment, primer extension products are characterized by liquid or gas chromatography, which may further include high performance liquid chromatography (HPLC), gas chromatography-mass spectrometry (GC-MS) or other well known chromatographic methodologies.

Any exmtDNA sequence or portion of an exmtDNA sequence that corresponds to the human mtDNA sequence of SEQ ID NO:2 or a portion thereof or several portions thereof may be useful in this embodiment of the invention. Examples of human exmtDNA sequences that are useful in this and other embodiments of the invention are disclosed in SEQ ID NO:1, in SEQ ID NO:3 and in SEQ ID NO:4.



Nucleotide positions at which exmtDNA and mtDNA differ are provided in Figure 4, in which are presented the identities of nucleotides at particular sequence positions in the human mtDNA genome (SEQ ID NO:2) and the corresponding positions in exmtDNA of SEQ ID NOS:1, 3 and 4, and in other specific exmtDNA regions corresponding to the human mtDNA encoded genes NADH dehydrogenase subunit 1 (ND1), NADH dehydrogenase subunit 2 (ND2), NADH dehydrogenase subunit 3 (ND3), NADH dehydrogenase subunit 4 (ND4), NADH dehydrogenase subunit 4L (ND4L), NADH dehydrogenase subunit 5 (ND5), NADH dehydrogenase subunit 6 (ND6), tRNA<sup>Lys</sup>, tRNA<sup>Ile</sup>, tRNA<sup>Gln</sup>, tRNA<sup>Met</sup>, tRNA<sup>Cys</sup>, tRNA<sup>Tyr</sup>, tRNA<sup>Asp</sup> or cytochrome c oxidase 3 (CO3). These exmtDNA nucleotide sequences that correspond to the human mtDNA sequence of SEQ ID NO:2 and that differ from SEQ ID NO:2 by the specific nucleotide substitution or deletion provided in Figure 4 for an indicated nucleotide position number (using the numbering system of Anderson et al., 1981 *Nature* 290:457) were determined as provided herein by sequence analysis of SEQ ID NOS:1, 3 or of SEQ ID NO:4 or of cloned PCR amplicons generated as described herein using oligonucleotide primer sets disclosed in Tables 1-3.

Portions of SEQ ID NO:2 that include nucleic acid sequences encoding the mitochondrial ETC enzymes cytochrome c oxidase 1 (CO 1), cytochrome c oxidase 2 (CO 2), ATP synthetase subunit 8 (ATPase 8) and ATP synthetase subunit 6 (ATPase 6) may be particularly useful, and in preferred embodiments of the invention these sequences comprise isolated nucleic acid molecules that have nucleotide sequences identical or complementary to corresponding nucleic acid sequences present in exmtDNA of SEQ ID NO:1 and/or SEQ ID NO:3 and/or SEQ ID NO:4. Portions of SEQ ID NO:2 that include nucleic acid sequences encoding the mitochondrial tRNAs, including mitochondrial isoleucyl, glutaminyl, methionyl, aspartyl, cysteinyl, tyrosinyl and lysyl tRNAs may also be particularly useful. Also particularly useful in preferred embodiments of the invention are sequences comprising isolated nucleic acid molecules that have nucleotide sequences identical or complementary to corresponding nucleic acid sequences present in exmtDNA of SEQ ID NO:1 and/or of SEQ ID NOS:3 and/or 4, and/or to corresponding regions of SEQ ID NO:2 having the specific nucleotide

substitutions or deletions identified at the indicated corresponding mtDNA nucleotide position in Figure 4, or any portion or fragment thereof.

Figure 4 shows nucleotides present in human mtDNA of SEQ ID NO:2 (Anderson et al., 1981 *Nature* 290:457) at the indicated nucleotide position number, and  
5 also shows single base substitutions or deletions, which substitutions and deletions are provided by the instant invention, at the corresponding nucleotide position number in human exmtDNA sequences. The "gene" column refers to the mtDNA encoded gene region in which the indicated mtDNA nucleotide position numbers are present: "COI", "COII" and COIII" refer, respectively, to cytochrome c oxidase subunits I (COX I), II  
10 (COX II) and III (COX III); "ATPase" refers to ATP synthetase (also commonly referred to as ATP synthase); "ND" refers to NADH dehydrogenase (also known as NADH:ubiquinone oxidoreductase or mitochondrial electron transport chain Complex I); and genes encoding tRNA specific for the amino acid indicated by the well known three letter code are appropriately labeled.

15 The "fragment" column in Figure 4 refers either (i) to cloned and sequenced exmtDNA containing the indicated nucleotide substitutions and deletions relative to the corresponding positions in human mtDNA, wherein "5.8 kb pg" refers to SEQ ID NO:1 and "519 bp pg" refers to SEQ ID NO:3; or alternatively, (ii) identifies oligonucleotide primer sets from Table 1 (COX I, II and III), Table 2 (ND1-4, 4L, 5 and  
20 6), or Table 3 (ATP synthetase subunits 8 and 6) that were used to PCR amplify cellular DNA as described herein, using a two-digit number with a letter code (A = alternate, i.e., an alternate primer complementary to an overlapping and/or adjacent region of human mtDNA [SEQ ID NO:2] relative to a first primer having the same two-digit number and orientation -- such alternates are included to provide means for PCR  
25 amplifying a particular human exmtDNA region containing one or more substitutions or deletions listed in Figure 4, where under certain conditions one member of the primer pair comprising the first primer and its alternate may be incapable of duplex formation with an exmtDNA target sequence due to an uncharacterized sequence variation, for example a mutation such as a point mutation, in the target sequence. Typically, an  
30 alternate primer will be designed to anneal to an exmtDNA target sequence that

commences at a nucleotide corresponding to a nucleotide position in human mtDNA situated within 1-100 nucleotides of the position where the first primer anneals. preferably within 1-50 nucleotides. more preferably within 1-30 nucleotides and still more preferably within 1-15 nucleotides. F = forward orientation; R =, reverse orientation.)

5

TABLE I  
OLIGONUCLEOTIDE PRIMERS CORRESPONDING TO CYTOCHROME  
C OXIDASE ENCODING MTDNA SEQUENCES

SEQ. ID. NO.	PRIMER	GENE	NUCLEOTIDE POSITION (5', light strand)	PRIMER SEQUENCE 5'→3'
14	11F	CO I	5864	GTCCAATGCTTCACTCAGCCA
15	11FA	CO I	5859	TTACAGTCCAATGCTTCACTC
16	11R	CO I	6177	TATGCGGGGAAACGCCAT
17	11RA	CO I	6180	TGTTTATGCGGGGAAACGC
18	12F	CO I	6138	GGCAACTGACTAGTTCCCCTA
19	12FA	CO I	6125	AATCGGAGGCTTTGGCAACTG
20	12R	CO I	6425	GTTTGGTATTGGGTTATGGCA
21	12RA	CO I	6422	TGGTATTGGGTTATGGCAGGG
22	13F	CO I	6383	GGCCATCAATTTCATCACAA
23	13FA	CO I	6358	TAGCAGGTGTCTCCTCTATCTT
24	13R	CO I	6697	ATACCTATGTATCCAAATGGTTCTT
25	13RA	CO I	6699	CCATACCTATGTATCCAAAIGGTTC
26	14F	CO I	6657	GGAATAATCTCCCATATTGTAACCT
27	14R	CO I	6945	CAGGCCACCTACGGTGAA
28	14RA	CO I	6947	GTCAGGCCACCTACGGTG
29	15F	CO I	6914	AGTGCTCTGAGCCCTAGGAT
30	15FA	CO I	6902	ATGATCTGCTGCAGTGCTCT
31	15R	CO I	7193	ATTCCGGATAGGCCGAGA
32	16F	CO I	7159	TCGGCGTAAATCTAACTTTCTT
33	16R	CO I	7451	GGGGTTCGATTCTTCCTT
34	16RA	CO I	7472	TTGGCTTGAAACCAGCTTT
35	21F	CO II	7546	TTGTCAAAGTTAAATTATAGGCTA
36	21FA	CO II	7548	GTCAAAGTTAAATTATAGGCTAAA
37	21R	CO II	7832	ACCTCGTCTGTTATGTAAAGGAT
38	21RA	CO II	7834	TGACCTCGTCTGTTATGTAAAGG
39	22F	CO II	7792	CGCCATCATCCTAGTCCTCA
40	22R	CO II	8050	ATGAGTGCAAGACGTCTTGTGAT
41	23F	CO II	8003	AATCGAGTAGTACTCCCGATTGA



SEQ. ID. NO.	PRIMER	GENE	NUCLEOTIDE POSITION (5', light strand)	PRIMER SEQUENCE 5'→3'
42	23FA	CO II	8007	GAGTAGTACTCCCGATTGAAGCC
43	23R	CO II	8286	GTTAGCTTTACAGTGGGCTCTAGA
44	23RA	CO II	8287	AGTTAGCTTTACAGTGGGCTCTAG
45	31F	CO III	9171	ACTTCTAGTAAGCCTCTACCTGCA
46	31FA	CO III	9173	TTCTAGTAAGCCTCTACCTGCACG
47	31R	CO III	9447	AGGTAATAAATAGGATTATCCCGTA
48	31RA	CO III	9443	AATAAATAGGATTATCCCGTATCGA
49	32F	CO III	9416	CCACACACCACCTGTCCA
50	32FA	CO III	9415	ACCACACACCACCTGTCCA
51	32R	CO III	9741	AAGGGAGACTCGAAGTACTCTGA
52	33F	CO III	9712	TGGGTCTCTATTTTACCCTCCTA
53	33FA	CO III	9698	TATTACAATTTTACTGGGTCTCT
54	33R	CO III	10010	ACTAGTTAATTGGAAGTTAACGGTA

TABLE 2  
OLIGONUCLEOTIDE PRIMERS CORRESPONDING TO NADH  
DEHYDROGENASE ENCODING MTDNA SEQUENCES

SEQ. ID. NO.	PRIMER	GENE	NUCLEOTIDE POSITION (5', light strand)	PRIMER LENGTH	PRIMER SEQUENCE 5'→3'
55	61F	ND1	3281	23	GAGGTTCAATTCCTCTTCTTAAC
56	61R	ND1	3631	22	TTGAGTAAACGGCTAGGCTAGA
57	62F	ND1	3589	20	CTGGTCAACCTCAACCTAGG
58	62R	ND1	3946	16	GGCCTGCGGCGTATTC
59	63F	ND1	3908	17	CCGAAGGGGAGTCCGAA
60	63R	ND1	4270	25	ATCAAAGTAACCTCTTTATCAGACA
61	71F	ND2	4447	20	TTGGTTATACCCTTCCCGTA
62	71R	ND2	4769	26	CTATTCCTAGTTTTATTGCTATAGCT
63	72F	ND2	4699	22	ACAATATACTCTCCGGACAATG
64	72R	ND2	5054	23	GAATGGTTATGTTAGGGTTGTAC
65	73F	ND2	4990	22	AGCTACGCAAAATCTTAGCATA
66	73R	ND2	5311	19	AGGGTGATGGTGGCTATGA
67	74F	ND2	5234	15	CCCGCTAACCGGCTT
68	74R	ND2	5536	18	AGGGCTTTGAAGGCTCTT
69	81F	ND4L	10433	25	TTTCGACTCATTAAATTATGATAAT
70	81R	ND4L	10782	25	CATGTCAGTGGTAGTAATATAATTG
71	82F	ND4	10718	22	CACATATGGCCTAGACTACGTA
72	82R	ND4	11060	23	ATAATTAAGGAGATTTGTAGGGA
73	83F	ND4	10999	19	CCAACGCCACTTATCCAGT
74	83R	ND4	11342	25	AAGCTATTGTGTAAGCTAGTCATAT
75	84F	ND4	11275	25	CTCACTAAACATTCTACTACTCACT
76	84R	ND4	11618	21	GTGGCTGATTGAAGAGTATGC
77	85F	ND4	11554	24	CCTATGAGGCATAATTATAACAAG
78	85R	ND4	11894	22	ACGTGGTTACTAGCACAGAGAG
79	86F	ND4	11834	19	TGACTTCTAGCAAGCCTCG
80	86R	ND4	12147	23	ACAATCTGATGTTTTGGTTAAAC
81	91F	ND6	14119	22	CTCATCCTAACCTACTCCTAA
82	91R	ND6	14442	19	GCGATGGCTATTGAGGAGT
83	92F	ND6	14384	20	GCTAACCCCACTAAACACT
84	92R	ND6	14693	20	TTCATATCATTGGTCGTGGT
85	75F	ND3	10007	25	TAGTACCGTTAACTTCCAATTAAC
86	75R	ND3	10430	24	TCATAATTTAATGAGTCGAAATCA
87	93F	ND5	12281	21	CAGCTATCCATTGGTCTTAGG
88	93R	ND5	12671	25	TATTTGAAGAACTGATTAATGTTTG
89	94F	ND5	12612	21	AGCATTGTTCTGTTACATGGTC
90	94R	ND5	12944	21	GGCTTGGATTAGCGTTTAGAA
91	95F	ND5	12881	20	TCATCCTCGCCTTAGCATGA
92	95R	ND5	13212	23	TTTTGATGTCATTTTGTGTAAGG
93	96F	ND5	13156	23	CAAACCTAACACTATGCTTAGG
94	96R	ND5	13519	20	ATGTTTGCGGTTTCGATGAT

SEQ. ID. NO.	PRIMER	GENE	NUCLEOTIDE POSITION (5', light strand)	PRIMER LENGTH	PRIMER SEQUENCE 5'→3'
95	97F	ND5	13458	20	CATTGGCAGCCTAGCATTAG
96	97R	ND5	13855	26	GATTTTATTTTAAGTTTGTTGGTTAG
97	98F	ND5	13795	19	AAACTCACAGCCCTCGCTG
98	98R	ND5	14124	24	TATGTGATTAGGAGTAGGGTTAGG

TABLE 3  
OLIGONUCLEOTIDE PRIMERS CORRESPONDING TO ATP SYNTHETASE SUBUNITS 8/6  
ENCODING MTDNA SEQUENCES

SEQ. ID. NO.	PRIMER	GENE	NUCLEOTIDE POSITION (5', light strand)	PRIMER LENGTH	PRIMER SEQUENCE 5'→3'
99	41F	ATPase 8	8292	23	GCCCACTGTAAAGCTAACTTAGC
100	41R	ATPase 8	8631	22	TAGTCGGTTGTTGATGAGATAT
101	41RA	ATPase 8	8632	17	AGTCGGTTGTTGATGAG
102	42F	ATPase 6	8572	23	GGCCTACCCGCCGCAGTACTGAT
103	42R	ATPase 6	8909	23	TGTAGGTGTGCCTTGTGGTAAGA
104	43F	ATPase 6	8866	22	ATTATAGGCTTTCGCTCTAAGA
105	43FA	ATPase 6	8806	21	CCAACCACCCAATATCTATA
106	43R	ATPase 6	9214	22	ATATGATAGGCATGTGATTGGT
107	26F	ATPase 8	8311	25	TAGCATTAAACCTTTTAAGTTAAAGA
108	26R	ATPase 8	8516	19	TCGTTCAATTTGGTTCTCA

5

As described herein, the present invention provides compositions and methods related to novel exmtDNA sequences that may differ from human mtDNA sequences at one or more nucleotide positions as disclosed in Figure 4, such as, for example, the exmtDNA sequences of SEQ ID NOS:1, 3 and 4. Details for obtaining

10 SEQ ID NO:1 are provided below in the Examples. Those having ordinary skill in the art can also readily obtain other isolated exmtDNA sequences using well known methodologies including those provided herein and in the cited references, and further including the use of the oligonucleotide primers provided in Tables 1-3 and in the Examples. Databases (e.g., GenBank, EMBL) and methods for nucleic acid sequence

15 analysis are also well known in the art, for example, similarity between two sequences may be readily determined using well known computer programs such as the BLAST algorithm (Altschul, *J. Mol. Biol.* 219:555-565, 1991; Henikoff and Henikoff, *Proc.*

*Natl. Acad. Sci. USA* 89:10915-10919, 1992). which is available at the NCBI website (<http://www.ncbi.nlm.nih.gov/cgi-bin/BLAST>). Default parameters may be used. Examples of other useful computer algorithms are those used in programs such as Align and FASTA, which may be accessed, for example, at the Genestream internet website of the Institut de Genetique Humaine, Montpellier, France (www2.igh.cnrs.fr/home.eng.html) and used with default parameters.

For instance, exmtDNA sequences present in SEQ ID NOS:3 or 4 that correspond to specific regions of SEQ ID NO:2 can be readily identified, and appropriate primers selected based on the sequences provided in Tables 1-3 and the exmtDNA nucleotide substitutions and deletions relative to the corresponding human mtDNA sequence (SEQ IDNO:2) provided in Figure 4. For example, SEQ ID NO:3 may be derived from, or detected in, a suitable biological sample using primer set 26 and/or primer set 41 as provided in Table 3, and methodologies essentially as provided in U.S. Patent No. 5,840,493 (and references cited therein), which is hereby incorporated by reference. As another example, such an approach applied to a biological sample comprising a cloned human peripheral blood leukocyte derived genomic DNA library (Clontech, Palo Alto, CA, catalog number HL1111j) probed with a nucleotide sequence corresponding to the region of SEQ ID NO:2 encoding ATP synthase subunit 8 may provide SEQ ID NO:4, portions of which may be useful in the subject invention compositions and methods. From the disclosure herein, as will be readily apparent to those having ordinary skill in the art, the nucleotide sequence from position 215-733 in SEQ ID NO:4 corresponds to SEQ ID NO:3, and both SEQ ID NOS:3 and 4 correspond to a portion of SEQ ID NO:2 that differs from SEQ ID NO:2 by having specific nucleotide substitutions disclosed in Figure 4. The nucleotide sequences in SEQ ID NO:4 from positions 1-214 and 734-1263, conversely, do not correspond to mtDNA sequences of SEQ ID NO:2 and further may represent novel nucleic acid sequences. Sequence analysis (*e.g.*, using online databases and algorithms as described above) of nucleotides 734-1263 suggests that nucleotides 963-1105 encode an exon of a human carboxypeptidase-N "L" subunit and that nucleotides 734-962 represent novel intron sequences within the carboxypeptidase-N gene, with nucleotides

1106-1263 representing additional novel (and probably non-coding) sequences as well. The exmtDNA sequence of nucleotides 215-733 thus may be present in human genomic DNA as a mtDNA-like pseudogene situated within intronic DNA.

In another particularly preferred embodiment of the invention, DNA in a biological sample containing exmtDNA and/or mtDNA is first amplified by methodologies well known in the art and described above, such that the amplification products may be used as templates in a method for quantifying the amount of exmtDNA and mtDNA present in the sample. Accordingly, it may be desirable to employ oligonucleotide primers that are complementary to target sequences that are identical in, and common to, mtDNA and exmtDNA, for example PCR amplification templates and primers prepared according to Fahy et al. (*Nucl. Acids Res.*, 25:3102, 1997) and Davis et al. (*Proc. Nat. Acad. Sci. USA* 94:4526, 1997; see also Hirano et al., *Proc. Nat. Acad. Sci. USA* 94:14894, 1997, and Wallace et al., *Proc. Nat. Acad. Sci. USA* 94:14900, 1997.)

Biological samples containing exmtDNA and mtDNA may comprise any tissue or cell preparation in which exmtDNA and mtDNA may be present. Biological samples may be provided by obtaining a blood sample, biopsy specimen, tissue explant, organ culture or any other tissue or cell preparation from a subject or a biological source. The subject or biological source may be a human or non-human animal, a primary cell culture or culture adapted cell line including but not limited to genetically engineered cell lines that may contain chromasomally integrated or episomal recombinant nucleic acid sequences, immortalized or immortalizable cell lines, somatic cell hybrid or cytoplasmic hybrid "cybrid" cell lines, differentiated or differentiable cell lines, transformed cell lines and the like. In certain preferred embodiments of the invention, the subject or biological source may be suspected of having or being at risk for having a disease associated with altered mitochondrial function, and in certain preferred embodiments of the invention the subject or biological source may be known to be free of a risk or presence of such as disease.

In certain other preferred embodiments where it is desirable to determine whether or not a subject or biological source falls within clinical parameters indicative

of Alzheimer's disease (AD). signs and symptoms of AD that are accepted by those skilled in the art may be used to so designate a subject or biological source, for example clinical signs referred to in McKhann et al. (*Neurology* 34:939, 1984, National Institute of Neurology, Communicative Disorders and Stroke and Alzheimer's Disease and  
5 Related Disorders Association Criteria of Probable AD, NINCDS-ADRDA) and references cited therein, or other means known in the art for diagnosing AD.

In certain aspects of the invention, biological samples containing mtDNA and exmtDNA may be obtained from the subject or biological source before and after contacting the subject or biological source with a candidate agent, for example  
10 to identify a candidate agent capable of effecting a change in the value of the ratio  $r$ , defined above, relative to the value of  $r$  before exposure of the subject or biological source to the agent.

In a most preferred embodiment of the invention, the biological sample containing mtDNA and exmtDNA may comprise a crude buffy coat fraction of whole  
15 blood, which is known in the art to comprise further a particulate fraction of whole blood enriched in white blood cells and platelets and substantially depleted of erythrocytes. Those familiar with the art will know how to prepare such a buffy coat fraction, which may be prepared by differential density sedimentation of blood components under defined conditions, including the use of density dependent separation  
20 media, or by other methods.

In another most preferred embodiment of the invention, the amount of exmtDNA and mtDNA in a biological sample may be quantified by first heating the sample in water to lyse cells contained therein, and then extracting cellular DNA from the lysed cells using an aqueous DNA extraction procedure. "Heating" may involve  
25 treating the cells for various times, typically 1-120 minutes, at a high temperature that is at least 80°C, preferably at least 90°C, more preferably at least 95°C and most preferably in a boiling water bath. Based on the compositions and methods disclosed in the present application, the ordinarily skilled artisan will be able to readily determine optimal times and temperatures for heating samples to practice the invention without  
30 undue experimentation. As used herein, an "aqueous DNA extraction" method refers to

preparation of DNA from such a boiled cell lysate without subjecting the lysate to sodium dodecylsulfate(SDS)/proteinase K treatments and/or without fractionating the lysate using a phenol-chloroform two-phase separation extraction step. Those skilled in the art will be familiar with various standard procedures for preparing and handling

5 DNA without the use of SDS/ proteinase K and/or phenol-chloroform.

According to certain embodiments of the invention, the particular cell type or tissue type from which a biological sample is obtained may influence qualitative or quantitative aspects of the exmtDNA and/or mtDNA contained therein relative to exmtDNA and/or mtDNA obtained from distinct cell or tissue types of a common

10 biological source. As described above, some diseases associated with altered mitochondrial function may manifest themselves in particular cell or tissue types. For example, AD is primarily a neurodegenerative disease that particularly effects changes in the central nervous system (CNS). It is therefore within the contemplation of the invention to quantify exmtDNA and mtDNA in biological samples from different cell

15 or tissue types as may render the advantages of the invention most useful for a particular disease associated with altered mitochondrial function, and the relevant cell or tissue types will be known to those familiar with such diseases.

In order to determine whether a mitochondrial alteration may contribute to a particular disease state, it may be useful to construct a model system for diagnostic

20 tests and for screening candidate therapeutic agents in which the nuclear genetic background may be held constant while the mitochondrial genome is modified. It is known in the art to deplete mitochondrial DNA from cultured cells to produce  $\rho^0$  cells, thereby preventing expression and replication of mitochondrial genes and inactivating mitochondrial function. See, for example, International Publication Number WO

25 95/26973, which is hereby incorporated by reference in its entirety, and references cited therein. It is further known in the art to repopulate such  $\rho^0$  cells with mitochondria derived from foreign cells in order to assess the contribution of the donor mitochondrial genotype to the respiratory phenotype of the recipient cells. Such cytoplasmic hybrid cells, containing genomic and mitochondrial DNAs of differing biological origins, are

30 known as cybrids.

"p<sup>0</sup> cells" are cells essentially completely depleted of mtDNA, and therefore have no functional mitochondrial respiration/ electron transport activity. Such absence of mitochondrial respiration may be established by demonstrating a lack of oxygen consumption by intact cells in the absence of glucose, and/or by demonstrating a lack of catalytic activity of electron transport chain enzyme complexes having subunits encoded by mtDNA, using methods well known in the art. (See, e.g., Miller et al., *J. Neurochem.* 67:1897-1907, 1996.) That cells have become p<sup>0</sup> cells may be further established by demonstrating that no mtDNA sequences are detectable within the cells. For example, using standard techniques well known to those familiar with the art, cellular mtDNA content may be measured using slot blot analysis of 1 µg total cellular DNA probed with a mtDNA-specific oligonucleotide probe radiolabeled with, e.g., <sup>32</sup>P to a specific activity ≥ 900 Ci/gm. Under these conditions p<sup>0</sup> cells yield no detectable hybridizing probe signal. Alternatively, any other method known in the art for detecting the presence of mtDNA in a sample may be used that provides comparable sensitivity.

"Mitochondrial DNA depleted" cells ("mtDNA depleted cells") are cells substantially but not completely depleted of functional mitochondria and/or mitochondrial DNA, by any method useful for this purpose. MtDNA depleted cells are preferably at least 80% depleted of mtDNA as measured using the slot blot assay described above for the determination of the presence of p<sup>0</sup> cells, and more preferably at least 90% depleted of mtDNA. Most preferably, mtDNA depleted cells are depleted of >95% of their mtDNA.

Mitochondria to be transferred to construct model systems in accordance with the present invention may be isolated from virtually any tissue or cell source. Cell cultures of all types may potentially be used, as may cells from any tissue. However, fibroblasts, brain tissue, myoblasts and platelets are preferred sources of donor mitochondria. Platelets are the most preferred, in part because of their ready abundance, and their lack of nuclear DNA. This preference is not meant to constitute a limitation on the range of cell types that may be used as donor sources.



For example, platelets may be isolated by an adaptation of the method of Chomyn (*Am. J. Hum. Genet.* 54:966-974, 1994). However, it is not necessary that this particular method be used. Other methods are easily substituted. For instance, if nucleated cells are used, cell enucleation and isolation of mitochondria isolation can be performed as described by Chomyn et al., *Mol. Cell. Biol.* 11:2236-2244, 1991. Human tissue from a subject suspected of having or being at risk for having a disease associated with altered mitochondrial function, or from a subject known to be free of a risk or presence of such a disease, may be the source of donor mitochondrial DNA.

After preparation of mitochondria by isolation of platelets or enucleation of donor cells, the mitochondria may be transplanted into  $\rho^0$  cells or mtDNA depleted cells using any known technique for introducing an organelle into a recipient cell, including but not limited to polyethylene glycol (PEG) mediated cell membrane fusion, cell membrane permeabilization, cell-cytoplasm fusion, virus mediated membrane fusion, liposome mediated fusion, particle mediated cellular uptake, microinjection or other methods known in the art. For example by way of illustration and not limitation, mitochondria donor cells ( $\sim 1 \times 10^7$ ) are suspended in calcium-free Dulbecco's modified Eagle (DME) medium and mixed with  $\rho^0$  cells ( $\sim 0.5 \times 10^6$ ) in a total volume of 2 ml for 5 minutes at room temperature. The cell mixture is pelleted by centrifugation and resuspended in 150  $\mu$ l PEG (PEG 1000, J.T. Baker, Inc., 50% w/v in DME). After 1.5 minutes, the cell suspension is diluted with normal  $\rho^0$  cell medium containing pyruvate, uridine and glucose, and maintained in tissue culture plates. Medium is replenished daily, and after one week medium lacking pyruvate and uridine is used to inhibit growth of unfused  $\rho^0$  cells. These or other methods known in the art may be employed to produce cytoplasmic hybrid, or "cybrid", cell lines.

As a non-limiting example, cybrid model systems may be useful for diagnosing a patient suspected of having or being at risk for a disease associated with altered mitochondrial function. According to this example, the patient's mitochondria are used to construct cybrid cells as described above. These cybrid cells may then be propagated in vitro and used to provide a biological sample for the determination of the ratio  $r$ , which can be compared to an  $r$  value calculated from samples of a control cybrid

cell line constructed with mitochondria from a subject known to be free of disease. Where it may be desirable to compare the influence upon  $r$  of mitochondria from different sources, both cybrid cell lines may be constructed from the same  $p^0$  cell line to provide a constant background environment. These and similar uses of model systems according to the invention for determining the risk for or presence of a disease associated with altered mitochondrial function will be appreciated by those familiar with the art and are within the scope and spirit of the invention.

As another non-limiting example, cybrid model systems may be useful for identifying agents suitable for treating a disease associated with altered mitochondrial function. According to this example, a cybrid cell line may be a biological source in which the ratio  $r$  is calculated as described above, before and after cybrid cells are contacted with a candidate agent for treating disease. Such a cybrid cell line may be used to screen candidate agents by identifying those agents capable of effecting a change in the value of  $r$  relative to the value of  $r$  before exposure to the agent. The present invention thus provides model systems for selecting therapeutic agents that may be suitable for the treatment of diseases associated with altered mitochondrial function. These and similar uses of model systems according to the invention for the screening and identification of agents that influence the ratio  $r$  defined above, will be appreciated by those familiar with the art and are within the scope and spirit of the invention.

In addition, although the present invention is directed primarily towards model systems for diseases in which the mitochondria have metabolic alterations, it is not so limited. Conceivably there are disorders wherein mitochondria contain structural or morphological defects or anomalies, and the model systems of the present invention are of value, for example, to find drugs that can address that particular aspect of the disease. Also, there are certain individuals that have or are suspected of having extraordinarily effective or efficient mitochondrial function, and the model systems of the present invention may be of value in studying such mitochondria. Moreover, it may be desirable to put known normal mitochondria into cell lines having disease characteristics, in order to rule out the possibility that mitochondrial alterations

contribute to pathogenesis. All of these and similar uses are within the scope of the present invention, and the use of the phrase "mitochondrial alteration" herein should not be construed to exclude such embodiments.

According to the present invention, a ratio  $r$  as defined herein is  
5 determined in a biological sample, for example by calculation following quantification of mtDNA and exmtDNA using a technique based on specific oligonucleotide hybridization to a target sequence. This hybridization may be optionally followed by target template directed extension, such as in primer extension assays described herein. For certain diseases associated with altered mitochondrial function, calculation of  $r$  may  
10 have diagnostic usefulness. For example, where other clinical indicators of a disease associated with altered mitochondrial function are known, values for  $r$  in subjects known to be free of a risk or presence of such disease based on the absence of these indicators may be determined to establish a control range for  $r$ . The ratio may also be calculated in biological samples obtained from subjects suspected of having or being at  
15 risk for having a disease associated with altered mitochondrial function, and compared to the control range of  $r$  values determined in disease free subjects. Those having familiarity with the art will appreciate that there may be any number of variations on the particular subjects, biological sources and bases for comparing  $r$  values that are useful beyond those that are expressly presented herein, and these additional uses are within  
20 the scope and spirit of the invention.

For instance, determination of  $r$  in may take the form of a diagnostic assay performed on whole blood collected from a subject by routine venous blood draw, on buffy coat cells prepared from blood or on biological samples that are other cells, organs or tissue from a subject. Alternatively, in certain situations it may be desirable  
25 to construct cybrid cell lines using mitochondria from either control subjects or subjects suspected of being at risk for a disease associated with altered mitochondrial function. Such cybrids may be used to determine  $r$  for diagnostic purposes, or as biological sources for screening assays to identify agents that may be suitable for treating disease based on their ability to change the  $r$  value obtained from treated cells. In one  
30 embodiment of this aspect of the invention, therapeutic agents or combinations of

agents that are tailored to effectively treat an individual patient's particular disease may be identified by routine screening of candidate agents on cybrid cells constructed with the patient's mitochondria.

The present invention provides compositions and methods that are useful  
5 in pharmacogenomics, for the classification and/or stratification of a subject or a patient population, for instance correlation of one or more traits in a subject with indicators of the responsiveness to, or efficacy of, a particular therapeutic treatment. In one aspect of the invention, measurement of  $r$  in a biological sample from a subject is combined with identification of the subject's apolipoprotein E (APOE) genotype to determine the risk  
10 for, or presence of, Alzheimer's disease (AD) in the subject. The apolipoprotein E type 4 allele (*APOE-ε4*) allele is a genetic susceptibility factor for sporadic AD and confers a two fold risk for AD (Corder et al., *Science* 261:921, 1993; see also "National Institute on Aging/Alzheimer's Association Working Group Consensus Statement," *Lancet* 347:1091, 1996 and references cited therein, all of which are hereby incorporated by  
15 reference in their entirety.). Accordingly, in a preferred embodiment of the invention, the method for determining the risk for or presence of AD in a subject by comparing  $r$  values will further comprise determining the APOE genotype of the subject suspected of being at risk for AD. By using the combination of the methods for determining  $r$ , as disclosed herein, and methods known in the art for determining APOE genotype, an  
20 enhanced ability to detect the relative risk for AD is provided by the instant invention along with other related advantages. Similarly, where APOE genotype and risk for AD are correlated, the present invention provides advantageous methods for identifying agents suitable for treating AD where such agents affect  $r$  in a biological source.

As described herein, determination of  $r$  may be used to stratify an AD  
25 patient population. Accordingly, in another preferred embodiment of the invention, determination of  $r$  in a biological sample from an AD subject may provide a useful correlative indicator for that subject. An AD subject so classified on the basis of an  $r$  value may then be monitored using AD clinical parameters referred to above, such that correlation between  $r$  value and any particular clinical score used to evaluate AD may  
30 be monitored. For example, stratification of an AD patient population according to  $r$

values may provide a useful marker with which to correlate the efficacy of any candidate therapeutic agent being used in AD subjects. In a further preferred embodiment of this aspect of the invention, determination of  $r$  in concert with determination of an AD subject's APOE genotype may also be useful. These and  
5 related advantages will be appreciated by those familiar with the art.

In another aspect, the invention provides exmtDNA sequences that may be useful in the detection of telomeric events that are related to diseases, including diseases associated with altered mitochondrial function, or in the identification of agents that are suitable for the treatment of such diseases. Telomeres are specialized DNA sequences  
10 found at the ends of eukaryotic chromosomes that are complexed *in vivo* with specific telomere-associated proteins. Typically, telomeres comprise tandemly repeated, highly conserved G-rich sequences (for example, the human repeat sequence is 5'-TTAGGG). The eukaryotic DNA polymerase  $\alpha$  is unable to replicate the very ends of linear DNA and thus every replication cycle results in a shortening of the telomeric ends in normal  
15 somatic cells. Without wishing to be bound by theory, this progressive shortening of telomeric length may contribute to molecular events that lead to cellular senescence (*e.g.*, a "mitotic clock"; see Harley et al., *Nature* 345:458-460, 1990; Greider, *Bioessays* 12:363-369, 1990, hereby incorporated by reference). According to this model, in order to avoid telomeric shortening germline cells employ the enzyme telomerase, a  
20 ribonucleoprotein that catalyzes the addition of the tandem repeat sequence to the ends of chromosomes (Kim et al., *Science* 226:2011-2015, 1994). Telomerase is also expressed in some tumors, suggesting that telomerase activation may be one means by which tumor cells develop from somatic cells (Shay et al., *Curr. Opin. Oncol.* 8:66-71, 1996).

25 Dynamic processes in the telomeric or distal end regions of chromosomes that involve specific nucleic acid sequences, and in particular that may involve particular nucleotide polymerase and nuclease activities, have thus been implicated in chromosomal events that may be related to cellular and molecular mechanisms of disease. See also, for example, Fossel, *J. Amer. Med. Assoc.* 279:1732  
30 (1998); LaBranche et al., *Nat. Genet.* 19:199 (1998); Shay, *Cancer J. Sci. Am.* 4:526

(1998): Nowak et al., *Cancer J. Sci. Am.* 4:148 (1998); Iwama et al., *Hum. Genet.* 102:397 (1998), all of which are hereby incorporated by reference. In one embodiment of the invention, nucleic acid sequences are provided that may be used to monitor telomeric events, including but not limited to telomerase activity. Accordingly, because  
5 telomeric structure may be related to cellular growth potential and/or senescence, as noted above by way of non-limiting theory, nucleic acid based determination of telomeric structure may provide effective means for the detection of related disease processes. The present invention provides identification of human exmtDNA sequences in human chromosomal telomeric regions, and other related advantages.

10 The following examples are offered by way of illustration, and not by way of limitation.

### EXAMPLES

#### EXAMPLE 1

##### CLONING OF EXMTDNA FROM PERIPHERAL 15 BLOOD LYMPHOCYTE GENOMIC DNA LIBRARY

Peripheral blood lymphocytes were separated from the peripheral blood mononuclear cell fraction of freshly drawn venous blood from healthy human volunteers and the DNA extracted by standard techniques. Plasmid isolation, production of competent cells, transformation and manipulations using cloning vectors  
20 were performed essentially as described (Sambrook et al., *Molecular Cloning, a Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989). The human lymphocyte DNA was partially digested with HindIII and inserted into the pBeloBAC11 vector (Genome Systems, Inc., St. Louis, MO with insert sizes ranging from 50 kb to over 240 kb to create a human genomic library. This library was  
25 screened by PCR using the following primers, which are complementary to mtDNA sequences in the CO1 encoding region of SEQ ID NO:2 but have single-base substitutions at the 3' end. Details of PCR reaction conditions are provided in US 5,565,323, which is hereby incorporated by reference, using 30 cycles instead of 25

cycles at 95°C and with a single reaction being performed instead of five separate reactions.

5'-CCTTACACCTAGCAGGTA SEQ ID NO:5

5'-ACGCCGATGAATATGATAGC SEQ ID NO:6

5 A single positive clone was identified having a genomic DNA insert that included exmtDNA and was expanded, with the DNA then being purified using Magnum KB-100 columns (Genome Systems, St. Louis, MO). Portions of the insert were amplified with internal PCR primers complementary to human mtDNA sequences (Anderson et al., *Nature* 290:456, 1981) and sequenced using Prism DyeDeoxy  
10 terminator chemistry (Perkin-Elmer, Foster City, CA) according to the manufacturer's instructions. Sequence information at the junctions of exmtDNA and adjoining non-mtDNA in the positive pBeloBAC clone was obtained using the BigDye Terminator cycle sequencing kit (Perkin-Elmer) according to the supplier's recommendations. The products of sequencing reactions were purified by ethanol precipitation or by using  
15 CentriSep spin columns (Princeton Separations, Princeton, NJ), then electrophoretically separated using an Applied Biosystems Model 373A DNA sequencing system (Applied Biosystems Division of Perkin-Elmer, Foster City, CA). Sequence Navigator software (Applied Biosystems) was used to analyze exmtDNA sequence data, and nucleotide insertions, deletions or substitutions were identified by comparing exmtDNA sequences  
20 to published human mtDNA sequence data. (SEQ ID NO:2, Anderson et al., *Nature* 290:456, 1981).

The insert contained the 5,840 base pair contiguous exmtDNA sequence of SEQ ID NO:1 (nucleotide positions 445-6284 in Figure 1), which corresponded to and exhibited 98% sequence homology with nucleotide positions 3914-9755 of the  
25 human mtDNA sequence of SEQ ID NO:2. There were 89 nucleotide positions at which substitutions or deletions (2) were detected, as indicated in Figure 4. The complete ~5.8 kb exmtDNA sequence is in a single reading frame relative to the corresponding region of the human mitochondrial genome (SEQ ID NO:2) with the exception of a two base pair deletion at nucleotide positions 8196-8197. Starting at the  
30 5' terminus (nt 445 in Fig. 1) and proceeding in the 3' direction, the exmtDNA

sequence includes DNA sequences corresponding, in order, to a truncated ND1 gene, complete ND2 and CO1 genes, a CO2 gene with the above noted two base pair deletion, complete ATP synthetase subunit 8 and ATP synthetase subunit 6 genes, interspersed tRNA genes (as indicated in Figure 4) and a truncated CO3 gene of the published  
5 human mtDNA sequence of SEQ ID NO:2 (Anderson et al., *Nature* 290:456, 1981). The non-mitochondrial DNA sequences on either side of the 5,840 base pair exmtDNA sequence (nucleotides 1-444 and 6285-6691) did not display homology to any nuclear DNA sequences listed in the GenBank database.



## EXAMPLE 2

## DETECTION OF EXMTDNA IN RHO-0 CELLS

In order to verify that presumptive exmtDNA sequences originated from nuclear and not mitochondrial DNA present in the DNA preparation from which the human genomic library was constructed, two established cell lines were depleted of mtDNA using ethidium bromide to generate  $\rho^0$  cells (Miller et al., *J. Neurochem.* 67:1897, 1996) and assayed for the presence of exmtDNA sequences. Briefly, p0118/5 and 064/5 SH-SY5Y neuroblastoma cells and 0A431 epidermal carcinoma cells were produced and maintained as described (Miller et al., 1996). Cells were harvested and DNA was extracted with DNazol (Molecular Research Center, Inc., Cincinnati, OH) according to the manufacturer's instructions. The recovered DNA was amplified by PCR and analyzed by primer extension assay using primers specific for a sequence within the mitochondrial COXI gene region containing the nucleotide at position 7146 of the mtDNA sequence (SEQ ID NO:2), which corresponds to a nucleotide substitution in the exmtDNA sequence. (Figure 4 and SEQ ID NO:1) PCR primers and reaction conditions and primer extension assays were as described in Fahy et al. (*Nucl. Acids Res.* 25:3102, 1997), which is hereby incorporated by reference in its entirety. Primer extension products corresponding to a region of the exmtDNA sequence of SEQ ID NO:1 and including a nucleotide corresponding to the guanosine residue at position 7146 or its complement were detected in  $\rho^0$  cells that contained no detectable mtDNA as described herein and in Miller et al. (*J. Neurochem* 67:1897, 1996).

## EXAMPLE 3

## ABSENCE OF DETECTABLE TRANSCRIPTS OF EXMTDNA SEQUENCES IN RT-PCR ASSAY

The reverse transcription-polymerase chain reaction (RT-PCR; Rappolee et al., *Science* 241:708, 1991; Chelly et al., *Nature* 333:858, 1988; Brenner et al., *BioTechniques* 7:1096, 1989) technique was employed to determine whether the novel 5.8 kb exmtDNA sequence cloned from a human genomic DNA library is transcribed. The buffy coat fraction of freshly drawn human venous blood was prepared using Accuspin devices (Sigma, St. Louis, MO) according to the manufacturer's instructions and total RNA was extracted from isolated buffy coat cells with Trizol reagent (GibcoBRL, Bethesda, MD) as recommended by the supplier. First strand cDNA was synthesized from poly-A<sup>+</sup> mRNA using the SuperScript™ preamplification system (GibcoBRL) with oligo(dT) as primer according to the manufacturer's instructions. PCR was conducted using this cDNA as template and primers complementary to portions of the mitochondrial cytochrome c oxidase subunit 2 (COII, COX2) mtDNA sequence (SEQ ID NO:2) as described in Fahy et al. (*Nucl. Ac. Res.* 25:3102, 1997). The corresponding region of exmtDNA (SEQ ID NO:1) contains nucleotide substitutions at positions 7650 and 7868, relative to mtDNA (Figure 4). Amplicons were purified and analyzed by the primer extension assay as described in Fahy et al. such that readily distinguishable products are predicted depending on whether or not the primer has hybridized to a target sequence adjacent to a sequence having the substitutions. Based on quantitative analysis of fluorescent band intensities of the primer extension products, mRNA encoding human COII gene products of mtDNA (SEQ ID NO:2) was detectable, but no mRNA encoding products from the corresponding exmtDNA region of SEQ ID NO:1 was detected, indicating that the exmtDNA sequence is not expressed.

## EXAMPLE 4

## PRIMER EXTENSION ASSAY TO QUANTIFY EXMTDNA AND MTDNA

A competitive primer extension approach was used to simultaneously detect both mtDNA and exmtDNA sequences with a fluorophor-labeled primer and a  
5 selected mix of deoxynucleotides (dNTPs) and dideoxynucleotides (ddNTPs). The exmtDNA and mtDNA compete as templates in the primer extension reaction and are distinguished by differential extension of the primer. Thus, to determine the relative quantities of defined portions of mtDNA (SEQ ID NO:2) and exmtDNA (SEQ ID NO:1) using the nucleotide substitution at position 7650 (Figure 4) with the nucleotide  
10 combination of dATP, dTTP and ddGTP, the primer is extended by one base when the template is mtDNA. When exmtDNA is present as template, the primer is extended by three bases. The proportion of exmtDNA in relation to mtDNA is estimated by comparing the ratio of fluorescence intensities of the gel-separated extension products with a standard curve generated from known mtDNA/exmtDNA mixtures. (Fahy et al.,  
15 *Nucl. Ac. Res.* 25:3102, 1997)

Thermo Sequenase™, dNTPs and ddNTPs were purchased from Amersham (Cleveland, OH). Calf intestine alkaline phosphatase and biochemical reagents were obtained from Boehringer Mannheim (Indianapolis, IN) and QIAquick PCR purification kits from Qiagen (Chatsworth, CA). Accuspin™ Tubes and  
20 HISTOPAQUE® 1077 were purchased from Sigma (St. Louis, MO) and EDTA vacutainers from Beckton Dickinson (San Jose, CA). *UITma*™ DNA polymerase, *AmpliTaq*® DNA polymerase and reagents for DNA synthesis were purchased from Perkin Elmer (Foster City, CA). Oligonucleotides were synthesized on an Applied Biosystems 394 DNA/RNA synthesizer (Perkin Elmer) using standard phosphoramidite  
25 chemistry. 5' Fluorescein-labeled oligonucleotide primers were obtained by using the 6-FAM Amidite reagent in the last step of the automated synthesis. Tritylated and fluorescein-labeled oligonucleotides were purified by reverse phase chromatography using an acetonitrile gradient in 0.1 M triethylammonium acetate, pH 6.8 running buffer. The oligonucleotides migrated as single bands on a 15% denaturing  
30 polyacrylamide gel. The homogeneity of the fluorescein-labeled oligonucleotides was

independently assessed by electrophoresis on an Applied Biosystems Model 373 Sequencing System.

After IRB approval and informed consent, fresh venous blood samples were drawn from 837 patients with clinical diagnosis of probable Alzheimer's disease (AD mean age =  $74.7 \pm 1.1$  years) and 191 controls ( $67 \pm 1$  years: cognitively normal age-matched N = 114; cortico-basal ganglionic degeneration N = 2; Pick's disease N = 1; Parkinson's disease N = 24; non-insulin dependent diabetes mellitus N = 29; insulin dependent diabetes mellitus N = 6; Leigh's syndrome, N = 2, Machado Joseph Disease N = 2; idiopathic renal glycosuria N = 1; progressive supranuclear palsy N = 1; sporadic amyotrophic lateral sclerosis N = 6; familial sporadic amyotrophic lateral sclerosis N = 1; familial Alzheimer's disease N = 2). AD patients met the National Institute of Neurological, Communicative Disorders and Stroke and Alzheimer's Diseases and Related Disorders Association (NINCDS-ADRDA) criteria of probable Alzheimer's disease (McKhann et al., *Neurology* 34:939, 1984).

Blood samples were collected in EDTA vacutainers and kept at 4°C for no more than 24 h. The platelet/white blood cell fraction was isolated with Accuspin™ Tubes (Sigma Diagnostics, St. Louis, MO) using the following procedure. Three ml of HISTOPAQUE® 1077 (Sigma) were added to the upper chamber of each Accuspin™ Tube and the device was centrifuged at 1000 g for 30 s. Two to three ml of blood were then introduced into the upper chamber and separated by centrifugation at 1000 g for 10 min at room temperature. After centrifugation, the plasma and white blood cell layers were transferred to a new tube and the white blood cells were sedimented by centrifugation at 7,000 g for 10 min. The white cell pellet was resuspended in 0.4 ml of a solution containing 0.9% sodium chloride/1mM EDTA and stored at -80°C until use.

Frozen white blood cells (0.2 mL) were thawed and were sedimented by centrifugation at 12,000 g for 5 min. The white cell pellet was washed twice with 0.6 ml of Dulbecco's Phosphate Buffered Saline (PBS; GibcoBRL, Bethesda, MD) and resuspended in 0.2 ml of water. The cells were lysed by incubation in a boiling water bath for 10 min. After cooling to room temperature, the cellular debris was sedimented by centrifugation at 14,000 g for 2 min. The supernatant was transferred to a new vial

and the approximate concentration of the crude DNA preparation was estimated from its A<sub>260</sub> absorbance. The DNA sample was stored at -80°C.

Primer extension reaction templates were prepared by PCR amplification of cellular DNA. The reactions were carried out in a total volume of 50 µl using the primer pair sets described below. Following amplification, the PCR products were analyzed by electrophoresis on a 0.8% agarose gel. Reactions for analyzing exmtDNA to mtDNA ratios contained ~1 µg of cellular DNA, 2.5 U of *AmpliTaq*® DNA polymerase, 20 pmol each of the light strand primer

5'-CATGCAGCGCAAGTAGGTCTACAAGAC-3' [SEQ ID NO:7]

and the heavy strand primer

5'-TGTTATGTAAAGGATGCGTAGGGATGG-3' [SEQ ID NO:8]

and 10 nmol of each dNTP in PCR buffer (10 mM Tris. HCl, pH 8.3, 50 mM KCl, 2 mM MgCl<sub>2</sub>). After an initial denaturation step at 95°C for 10 s in a Gene Amp PCR System 9600 thermal cycler (Perkin Elmer), amplification was carried out for 30 cycles under the following conditions: 95°C for 1 min, 60°C for 1 min, 72°C for 1 min. After the last cycle, reaction tubes were kept at 72°C for 4 min to ensure extension of incomplete strands to the full length 255 base pair size product.

Residual nucleotides that persisted after the PCR reaction were dephosphorylated by adding 1 unit of calf intestine alkaline phosphatase (CAP) in 5 µL of 10X CAP buffer (100 mM Tris. HCl, pH 8.3, 10 mM MgCl<sub>2</sub>, 10 mM ZnCl<sub>2</sub>) to the PCR reaction mixture and incubating for 30 min at 37°C in thermal cycler. Then 1.1 µL of 0.25 M EDTA, pH 8.0 was added and the alkaline phosphatase was denatured at 75°C for 10 min.

Double-stranded PCR products were separated from primers, nucleosides and enzymes using QIAquick™ columns (Qiagen, Chatsworth, CA) and the buffers provided by the manufacturer. Thus, 250 µl of buffer PB were added to the PCR

reaction mixture and mixed. A Qiaquick™ spin column (Qiagen) was placed in a 2 ml collection tube and the sample was loaded. The sample was centrifuged for 30-60 s at 14000 g and the flow through was discarded. The adsorbed PCR product was washed with 750 µl of buffer PE, and eluted with 50 µl of 10 mM Tris. HCl, pH 8.5. The  
5 purified product solution was dried in a Savant SpeedVac Concentrator and then reconstituted in 20 µl of water.

The fluorescein-labeled primer for analysis of the AD-associated mutation at nucleotide position 7650 was

10                   TATGAGGGCGTGATCATGAAAG                   [SEQ ID NO:9]

using dATP and dTTP plus dideoxyGTP (ddGTP) to generate primer extension products of 23 and 25 nucleotides in length from mtDNA and exmtDNA templates, respectively.

15                   Stock solutions of each dNTP and ddNTP were prepared by mixing equimolar amounts of the nucleotides with MgCl<sub>2</sub> and diluting the mixture to the desired concentration with 10 mM Tris, 1 mM EDTA, pH 8.0 (TE). The fluorescein-labeled primers were diluted in TE to provide final stock concentrations of 40 fmol/µl. One µl of the purified PCR-amplified DNA fragment was used as template for each  
20 assay.

Primer extension reactions were performed in a total volume of 8 µL. The thermostable enzyme, *UITma*™ DNA polymerase (Perkin-Elmer, Foster City, CA) was used in primer extension assays for analyzing exmtDNA to mtDNA ratios. The reactions contained template, 20 fmol fluorescein-labeled primer, 400 µM ddNTPs/25  
25 µM dNTPs of the appropriate nucleotide combination and 0.6 unit of enzyme in buffer containing 10 mM Tris-HCl, pH 8.8, 10 mM KCl, 0.002% Tween 20, 2 mM MgCl<sub>2</sub>. Each set of primer extension assays included control reactions with mtDNA and exmtDNA templates.

The products of the primer extension reaction were analyzed on an ABI  
30 373 Sequencer using a 12% denaturing polyacrylamide gel and Tris borate/EDTA as

running buffer. Prior to electrophoresis, the samples in loading dye were denatured for 3 min at 85°C. Three µl aliquots of the standards (primer with no added template, reaction products from control DNA templates) and each unknown reaction mixture were then loaded and electrophoresed according the manufacturer's instructions.

- 5 Fluorescent band intensities associated with the primer extension products were estimated by the GENESCAN™ 672 software program (Perkin Elmer, Applied Biosystems Division). Quantitative analysis was carried out by correlating the fluorescent band intensities of mtDNA and exmtDNA-derived extension products from unknown samples with those from mtDNA and exmtDNA control templates.

## EXAMPLE 5

PRIMER EXTENSION ASSAY USING DNA  
ISOLATED FROM AFFINITY ISOLATED MITOCHONDRIA

Mitochondria were immunoaffinity isolated from cultured cells and then  
5 DNA was extracted from the isolated organelles, to determine whether detectable  
exmtDNA was present. Normal SH-SY5Y neuroblastoma cells, and normal and  $\rho^0$   
A431 epidermal carcinoma cells were produced and maintained as described (Miller  
et al., *J. Neurochem.* 67:1897, 1996). Cells were harvested by scraping in MSB (0.21  
M mannitol, 0.07 M sucrose, 0.05 M Tris-HCl, 0.01 M EDTA, pH 7.4;  $10^7$  cells/ml)  
10 and lysed by three freeze-thaw cycles. Cellular debris was removed by centrifugation at  
1000 x g for 5 min. The mitochondria enriched supernatants were used for subsequent  
immunopurification steps. The buffy coat fraction of whole blood containing white  
blood cells and platelets as described above was also prepared and lysed by freeze-thaw  
as was done with the cultured cells, to produce a mitochondrial fraction for  
15 immunopurification.

Monoclonal anti-mitochondrial antibody MAB 1273 (Chemicon  
International, Temecula, CA) was added to the mitochondria enriched fractions for 2 hr.  
Antibody-mitochondria complexes were isolated using a secondary antibody bound to  
magnetic beads (Dynal Inc., Great Neck, NY) according to the manufacturer's  
20 instructions. After extensive washing of the bead-bound antibody-mitochondria  
complexes with PBS/0.1% BSA, mtDNA was extracted from the complexes using  
DNAzol reagent (Molecular Research Center, Inc., Cincinnati, OH) according to the  
manufacturer's recommendations. Quantification of mtDNA and exmtDNA was  
performed using the competitive primer extension assay described above in Example 4.  
25 No DNA could be amplified from the DNA extracted from immunopurified  
mitochondria from the  $\rho^0$  A431 cell line, consistent with the depletion of mtDNA that is  
characteristic of the  $\rho^0$  state. Primer extension results indicated that no exmtDNA  
sequences were present in the DNA extracted from mitochondria of blood buffy coat  
cells, SH-SY5Y cells or A431 cells, confirming that exmtDNA sequences are of  
30 extramitochondrial origin.



## EXAMPLE 6

## DETERMINATION OF APOE GENOTYPE BY PRIMER EXTENSION ASSAY

Primer extension assay procedures, essentially as described above in Example 4 but with the modifications indicated here, were used to determine APOE genotype in a panel of 837 clinically diagnosed AD individuals and 191 control subjects (cognitively normal age matched individuals, patients with non-insulin dependent diabetes mellitus (NIDDM) and neurological controls). The APOE allele distribution in the AD population and controls in the study is shown in Table 4.

TABLE 4

APOE GENOTYPE DISTRIBUTION

<i>APOE</i> allele	Control (N = 191)	AD (N = 837)
2/2	1	
2/3	16	33
2/4	3	22
3/3	112	295
3/4	56	403
4/4	3	84

DNA isolation and preparation of primer extension reaction templates by PCR were essentially as described above in Example 4, except that for *APOE* analysis, a modification of the protocol of Livak and Hainer (1994) was used. Thus, for template preparation by PCR each reaction contained ~1 µg of cellular DNA, 2.5 U of *AmpliTaq*® DNA polymerase, 20 pmol each of the forward primer

5'-GGCACGGCTGTCCAAGG-3'

[SEQ ID NO:10]

20 and the reverse primer

5'-CCCGGCCTGGTACACTG-3'

[SEQ ID NO:11]

and 10 nmol of each dNTP in PCR buffer (10 mM Tris. HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl<sub>2</sub> supplemented with 5% DMSO). After an initial denaturation step at 95°C for 10 s in a Gene Amp PCR System 9600 thermal cycler (Perkin Elmer), amplification was carried out for 25 cycles under the following conditions: 95°C for 1 min, 55°C for 1 min, 72°C for 1 min. The PCR product obtained after amplification was 226 base pairs in length.

The fluorescein-labeled primers and nucleotide combinations for primer extension analysis of Apo-E alleles are shown in Table 5.

TABLE 5

Gene	Primer Sequence (5'→3')	Size	Primer Extension Product Sizes	dNTP	ddNTP
APOE Codon 112	GCGGACATGGAGGACGTG SEQ ID NO:12	18	19, 20	T	G, C
APOE Codon 158	CGATGCCGATGACCTGCAGAAG SEQ ID NO:13	22	23, 24	T	G, C

Thermo Sequenase™ (Amersham, Cleveland, OH) DNA polymerase catalyzed reactions for Apo-E allele analysis were performed with 20 fmol fluorescein-labeled primer, 25 μM each of the appropriate ddNTP/dNTP combination and 0.64 unit of enzyme in buffer containing 10 mM Tris-HCl, pH 9.5, 5 mM KCl, 0.002% Tween 20, 2 mM MgCl<sub>2</sub>. Each set of primer extension assays included Apo-E allelic DNA controls. After an initial denaturation step at 95°C for 2 min, the reaction conditions comprised 20 cycles of 95°C for 20 s and 55°C for 40 s. The samples were concentrated to ~1 μl by heating open reaction tubes at 94°C for 7 min. After the concentration step, 8 μl of loading dye (0.5% blue dextran in 83% formamide/8.3 mM EDTA, pH 8.0) was added.

The products of the primer extension reaction were analyzed as described above in Example 4. The Apo-E allele composition of unknown DNA

samples was deduced by comparing the electrophoretic pattern of primer extension products with those from Apo-E allele DNA standards. APOE genotype distribution is shown in Table 4.

#### EXAMPLE 7

##### 5 CORRELATION OF EXMTDNA:MTDNA RATIO WITH RISK FOR AD

Primer extension assays as described in Example 4 were used to quantify the amounts of mtDNA and exmtDNA in blood samples from a panel of 837 clinically diagnosed AD individuals and 191 control subjects (cognitively normal age matched individuals, patients with non-insulin dependent diabetes mellitus (NIDDM) and  
10 neurological controls). For each subject, a ratio  $r$  was calculated using the formula:

$$r = x/(x + y)$$

wherein

15  $x$  is the amount of exmtDNA in a sample, and  
 $y$  is the amount of mtDNA in the sample.

The values for  $r$  were multiplied by 100 to give ranges of values according to which the patient population was stratified as shown in Table 6. Within each stratified range, the ratio of the number of subjects diagnosed with AD (McKhann et al., *Neurology* 34:939, 1984) to the number of subjects not having AD was  
20 determined, showing a correlation of increasing  $r$  value with increasing risk for AD (Table 6).

TABLE 6

Relative Risk for AD: by $r$ Value	
$r \times 100$	# of AD subjects: # of Non-AD subjects
>15	1.42
>20	1.8
>25	2.2
>30	2.0

## EXAMPLE 8

INCREASED PREDICTIVE VALUE OF EXMTDNA:MTDNA RATIO IN SUBJECTS HAVING  
AT LEAST ONE APOE4 ALLELE

Determination of the value  $r$  according to Example 7 was combined with  
 5 determination of APOE genotype according to Example 6 using the patient populations  
 characterized in Examples 6 and 7 to ascertain the relationship of APOE genotype to  
 relative risk for AD, and to demonstrate the increased correlative value of APOE  
 genotype combined with increasing  $r$  values with relative risk for AD. The relative risk  
 for AD as a function of APOE genotype using the APOE alleles 2, 3 or 4 alone is  
 10 shown in the bottom line of Table 7. The remainder of Table 7 shows the relative risk  
 of AD as a function of APOE genotype and  $r$  value when subjects are stratified  
 according to  $r$  as in Example 7. The relative risk for AD increases as a function of  
 increasing  $r$ . In particular individuals with one APOE4 allele, and especially subjects  
 who are homozygous for the APOE4 allele, exhibit increasing risk for AD as a function  
 15 of increasing  $r$ , as shown in Table 7.

TABLE 7

Relative Risk for AD: by $r$ value and ApoE Genotype (# AD: # Non-AD) ApoE Genotype					
$r \times 100$	e2/3	e3/3	one e4 allele	e3/4	e4/4
>15	1.5	1.5	2.2	2.1	7.1
>20	2.0	1.9	3.7	3.1	9.0
>25	1.7	1.8	5.3	3.0	>9.0
>30	2.5	2.0	4.6	3.0	>9.0
independent of $r$	0.5	0.6	1.9	1.7	6.3

## EXAMPLE 9

IDENTIFICATION OF AGENTS THAT ALTER THE VALUE OF  $r$ 

In this example, an agent suitable for treating AD is identified based on its ability to lower the value of  $r$  as defined above. A blood sample is obtained from a  
5 patient diagnosed with AD (McKhann et al., *Neurology* 34:939, 1984) and a ratio  $r$  is calculated using the formula:

$$r = x/(x + y)$$

10 wherein

$x$  is the amount of exmtDNA in a sample, and

$y$  is the amount of mtDNA in the sample, where  $x$  and  $y$  are determined using the primer extension assay as described in Example 4. The candidate agent is then administered to the patient in a quantity and for a time sufficient to impart a  
15 therapeutically beneficial effect, and blood samples are periodically collected and processed using the primer extension assay as described in Example 4 to monitor alterations in the value  $r$ . Candidate agents are subjected to preliminary characterization for toxicity, bioavailability and modes of delivery prior to administration to a patient. An agent is selected that causes a reduction in the value for  $r$ , signifying an effect on the  
20 altered mitochondrial function in the patient that may underlie the differential extractability of exmtDNA and mtDNA that contributes to  $r$  values associated with increased risk for AD.

## EXAMPLE 10

25

## DETECTION OF EXMTDNA SEQUENCES IN TELOMERES

In this example, fluorescence *in situ* hybridization (FISH) is used to localize nucleic acid sequences that are present in SEQ ID NO:1 to the telomeric region of at least one human chromosome. The methods of Cannizzaro et al. (*Methods Mol. Biol.* 75:313, 1997) and references cited therein, all of which are incorporated hereby in  
30 their entireties, are used to conduct FISH. Human cell lines are grown to confluence on

coverslips and then fixed and permeabilized for FISH analysis of metaphase chromosomes. The cloned pBeloBAC insert containing SEQ ID NO:1 as described above is excised from the plasmid vector and fluorescein labeled as described, and used to probe the fixed and permeabilized cells. Metaphase cells are evaluated by  
5 fluorescent laser scanning confocal microscopy and subchromosomal localization of the fluorescent probe to telomeric regions is observed. Cytological methods are used to prepare metaphase spreads of human peripheral blood leukocytes as described in the cited references, and the chromosomes are doubly labeled using suitable reporter moieties for discerning two signals, one being quinacrine to identify each chromosome  
10 by its characteristic banding pattern and the other being the labeled SEQ ID NO:1 insert, to correlate a particular chromosome with telomeric hybridization of the insert. The nucleic acid insert of SEQ ID NO:1 is next fragmented with restriction endonucleases, and the separated fragments are labeled to generate a panel of probes representing distinct sequence portions of SEQ ID NO:1, to determine whether the  
15 portion of the insert hybridizing to telomeres corresponds to an exmtDNA sequence or a non-exmtDNA sequence present in the insert as it was cloned from the human genomic library, as described above in Example 1.

#### EXAMPLE 11

##### 20 DETECTION OF EXMTDNA SEQUENCES IN TELOMERES BY RADIATION HYBRID MAPPING

This example describes radiation hybrid (RH) screening by PCR using the Stanford G3 panel (Research Genetics, Huntsville, AL) to positionally map exmtDNA sequences in the human genome. Localization of the exmtDNA sequences within the nuclear DNA was accomplished by radiation hybrid (RH) mapping according to  
25 Boehnke *et al.* (1991 *Am. J. Hum. Genet.* 49:1174) and Walter *et al.* (1994 *Nat. Genet.* 7:22). The Stanford G3 RH panel consists of more than eighty whole-genome radiation hamster-human hybrids and 14,000 characterized chromosomal markers available at the Stanford Human Genome Center (SHGC; <http://www-shgc.stanford.edu>). Eighty-three hybrid clones of the G3 RH panel, plus both parental human and hamster control cells,

were screened with PCR primers homologous to a portion of the CO III gene region within the exmtDNA sequence. PCR primers were designed to select for the exmtDNA sequence and contained nucleotide changes at positions corresponding to nucleotides 9325, 9329, 9335, 9540, 9545, and 9548 of SEQ ID NO:2:

5

L-strand primer: 5'- TCCACTCCACAACCCTCCTT SEQ ID NO:109

H-strand primer: 5'- GGGCCAGTGTCCCCTAG SEQ ID NO:110

10 Human and hamster genomic DNAs (Research Genetics, Huntsville, AL) were used to confirm specific amplification of a 223 bp fragment (exmtDNA marker) using AmpliTaq® DNA polymerase (Perkin-Elmer, Norwalk, CT) and standard PCR conditions as described above. Analysis was performed on the RH Server operated by the Stanford Human Genome Center (SHGC; <http://www-shgc.stanford.edu>, Palo Alto, CA). The exmtDNA marker was linked to more than 5,000 framework markers and EST's that have been characterized by the SHGC and by Genethon ([http://www.genethon.fr/genethon\\_en.html](http://www.genethon.fr/genethon_en.html)).

15 An amplification product of the expected 223 bp size was obtained from hybrid clone numbers 3, 11, 33, 40, 49, 50, 59, 64, 73, and 75, as well as from the human control cells, and not from the hamster control cells. The closest SHGC markers identified were SHGC-57364 with positive hybrid clone numbers 3, 11, 33, 40, 49, 50, 59, 64, 73, and 75; SHGC-4723 with positive hybrid clone numbers 11, 33, 49, 50, 51, 59, 64, 68, 73, and 75; and SHGC-428 with positive hybrid clone numbers 9, 11, 40, 49, 50, 51, 59, 64, 68, 73, and 75. All three markers were localized to a very distal portion at or near the telomeric region of chromosome 1, at distances from the exmtDNA of 0, 26, and 32 centirays (cR; one cR corresponds roughly to 26 kb), which is roughly equivalent to 0, 676, and 832 kb, respectively (Figure 5). SHGC-4723 and SHGC-428 are ordered markers linked to Bin#1 of chromosome 1 at GDB Locus D1S2565 and D1S243, respectively. SHGC-57364 is a positional marker, also linked to Bin#1 of chromosome 1, that has been oriented at a distance of 26 cR from SHGC-4723 (LOD

20

25

30

7.11). Interestingly, this marker (GenBank Accession No. T10998) originated from a 230 nucleotide sequence with partial similarity to the mitochondrial gene for cytochrome *c* oxidase subunit 2. It is therefore conceivable that the PCR primers used for identifying this marker in the clones where it was found may also have amplified the exmtDNA sequence. However, independent confirmation of the exmtDNA co-localization was provided by the specific CO III primer set described herein. These data demonstrate unambiguously the nuclear chromosomal location of this exmtDNA and its presence as a single copy sequence.

10

## EXAMPLE 12

## DETECTION OF EXMTDNA BY SOUTHERN BLOT HYBRIDIZATION ANALYSIS

The nucleotide substitution A→C at position 6266 in the portion of the exmtDNA sequence corresponding to the CO I gene (Fig. 4) creates a six-base pair *Nae* I restriction recognition site which is not present in human wild type mtDNA. In this example, this difference is exploited to distinguish mtDNA from the exmtDNA sequence by Southern blot analysis of human cellular DNA isolated from white blood cells (Figure 6a) or from p<sup>0</sup> cells (Fig. 6b), or of mtDNA from isolated human brain mitochondria (Fig. 6b).

DNA from two normal control individuals was restricted with *Sac* I, which cleaved at nucleotide position 9647 in the CO II gene of both the mtDNA and exmtDNA, and with *Nae* I, which cleaved at nucleotide position 932 in the 12S rRNA gene of mtDNA and nucleotide position 6266 in the exmtDNA. A wild type mitochondrial ATPase 8 probe detected the resulting 8715 bp mtDNA and 3381 bp exmtDNA fragments. The signal intensities of the two DNA fragments indicated their relative abundance in the cellular DNA preparation, suggesting a low copy number or possibly a single copy of this exmtDNA within the genomic DNA.

Human brain samples were obtained post mortem as soon as possible after death. The biological samples were frozen and stored at -80°C, and shipped on dry ice to analysis facilities. Mitochondria were purified from human brain tissue and mtDNA



was extracted as described by Mecocci *et al.* (1994 *Ann. Neurol.* 36:747-751). Briefly, human parietal cortex tissue was homogenized in an isotonic solution (0.05 M Tris-HCl, pH 8.0, 0.21 M mannitol, 0.07 M sucrose, 3 mM calcium chloride) using a glass dounce homogenizer and a round bottom glass rod with 0.15  $\mu$ m clearance. EDTA concentration was brought to 0.02 M and the homogenate was centrifuged at 1,500 g for 20 min at 4°C. The supernatant was transferred to a new tube, centrifuged again at 1,500 g for 20 min at 4°C, then transferred to a new tube and centrifuged at 17,000 g for 20 min at 4°C. The pellet consisting of mitochondria was resuspended in 40 mM Tris-HCl, pH 8.0, 10 mM magnesium chloride, containing DNase I and RNase A (Roche Molecular Biochemicals, Inc., Indianapolis, IN) at concentrations of 40 units/ml and 0.02 mg/ml, respectively, and incubated for 45 min at 37°C. The organelle suspension was diluted to 20 ml with isotonic dilution buffer (0.05 M Tris-HCl, pH 8.0, 0.21 M mannitol, 0.07 M sucrose, 0.01 M EDTA) and centrifuged at 17,000 g for 20 min at 20°C. The pellet was resuspended in isotonic dilution buffer, combined with an equal volume of lysis buffer (0.01 M Tris-HCl, pH 8.0, 0.1 M NaCl, 1 mM EDTA, 0.5% SDS) containing 400  $\mu$ g/ml Proteinase K (Roche Molecular Biochemicals, Inc., Indianapolis, IN) and incubated for 3 hr at 37°C. DNA was purified by three phenol extractions followed by three chloroform:isoamyl alcohol (24:1) extractions and ethanol precipitation. The DNA was resuspended in buffer (10 mM Tris-HCl, pH 7.4, 1 mM EDTA) and its concentration was determined by UV absorbance at 260 nm.

Total DNA isolated from white blood cells with DNAzol reagent, DNA isolated from p<sup>0</sup> 118/5 SH-SY5Y cells with DNAzol reagent, and DNA extracted from purified human brain mitochondria as described above were treated with *Nae* I and *Sac* I restriction endonucleases (New England Biolabs, Beverly, MA), separated by horizontal agarose gel electrophoresis, and blotted onto Nytran Plus membranes (Schleicher & Schuell, Keene, NH). Membranes were probed with a ~200 bp fragment of the wildtype ATP synthase subunit 8 (ATPase 8) gene that had been generated by PCR amplification of a DNA preparation from white blood cells using the following oligonucleotide primers:

L-strand primer: 5'-TAGCATTAACCTTTTAAGTTAAAGA SEQ ID NO:111

H-strand primer: 5'-TCGTTTCATTTGGTTCTCA SEQ ID NO:112

5 The probe was labeled with  $^{32}\text{P}$ -dATP (Amersham, Arlington Heights, IL) and the random priming method (Prime-a-Gene™ Labeling System, Promega, Madison, WI) according to the suppliers' recommendations. The blot was incubated with approximately 50 million cpm (5-10 ng) of probe for 16 h at 42°C in hybridization solution (6X SSC, 50% formamide, 1X Denhardt's solution, 1% SDS and 100 µg/ml salmon sperm DNA). Blots were washed twice for 5 min in 6X SSC, 0.1% SDS at room temperature, once for 30 min in 2X SSC, 0.1% SDS at 42°C, and once for 30 min in 0.2X SSC, 0.1% SDS at 42°C, and exposed to XAR 5 film (Kodak) for 1 to 4 days and developed.

15 Figure 6a shows hybridization of the ATPase 8 probe to total DNA isolated with DNazol reagent from white blood cells of two control individuals (A, B), and Figure 6b shows probe hybridization to DNA isolated from  $\rho^0$  SH-SY5Y cells with DNazol reagent and from purified human brain mitochondria with Proteinase K digestion, as described above. Approximately 10 µg of each DNA sample was restricted with *Nae* I and *Sac* I, and probed with the PCR amplified ATPase 8 probe. 20 Wildtype mtDNA carries a *Nae* I site at nucleotide position 932 which yields a 8,715 bp restriction fragment in combination with *Sac* I restriction at nucleotide position 9647. *Nae* I recognized a restriction site present in the exmtDNA sequence, which was created by the nucleotide substitution at position 6266 (Fig. 4) and was not present in the wildtype mtDNA. By cleavage of the exmtDNA at this site and at nucleotide position 25 9647 by *Sac* I, a 3,381 bp restriction fragment was created. An unlabeled aliquot (~4 ng) of the 200 bp ATPase 8 probe (Fig. 6a, probe control) showed an appropriate hybridization signal. Exposure times of the blots to X-ray film were (Fig. 6a) 24 hr and (Fig.6b) 4 d.

To provide additional support for an extramitochondrial (e.g., nuclear) 30 origin of the 5.8 kb exmtDNA sequence of SEQ ID NO:1, DNA from human  $\rho^0$  SH-

SY5Y cells (*i.e.*,  $\rho^0$  cells depleted of mtDNA as described in Miller et al., 1996 *J. Neurochem.* 67:1897) was restricted with *Sac* I and *Nae* I and hybridized with the wildtype mitochondrial ATPase 8 probe, which detected the resulting 3381 bp exmtDNA fragment (Figure 6b). To further demonstrate that the exmtDNA sequence  
5 does not reside within mitochondria, DNA was extracted from purified mitochondria derived from a human parietal cortex brain sample and digested with *Sac* I and *Nae* I, as described above. Following Southern blot hybridization with the wildtype mitochondrial ATPase 8 probe, an 8715 bp mtDNA fragment was detected but no detectable 3381 bp exmtDNA fragment was present (Figure 6b).

10

## EXAMPLE 13

## DETECTION OF TELOMERIC CHANGES USING EXMTDNA SEQUENCES

The nucleotide sequences of the 5.8 exmtDNA (SEQ ID NO:1), and its flanking regions, are used to design nucleic acid probes that specifically detect nucleic acids  
15 corresponding to the telomere region located at one end of human chromosome 1. Methods utilizing such probes are used to determine the degree of telomeric shortening of chromosome 1.

A synthetic oligonucleotide comprising a junction sequence of the exmtDNA is labeled and used as a probe of nuclear DNA in Southern blot analysis under stringent  
20 conditions, or as an unlabeled primer for PCR amplification of the telomeric region of chromosome 1. In PCR amplifications, the second primer is labeled and has a sequence specific for the telomere repeat sequence (5'-TTAGGG in humans); such primers are commercially available from Roche Molecular Biochemicals (Indianapolis, IN). The labeled PCR products are detected and their length assayed according to methods  
25 known in the art (*e.g.*, polyacrylamide gel electrophoresis (PAGE) with appropriate molecular weight markers). While not wishing to be bound by any theory, somatic cells from more aged individuals may have fewer numbers of telomeric repeats on the ends of their chromosomes than those from less aged individuals and, accordingly, the lengths of the PCR amplification products in biological samples derived from these

more aged individuals are shorter than those in biological samples derived from younger individuals. In like fashion, because some tumor cells may reactivate the enzyme (telomerase) that promotes the addition of telomeric repeats in germline (but not, typically, in somatic) cells, the lengths of the PCR amplification products in  
5 biological samples derived from a tumor are longer than those in biological samples derived from healthy individuals.

From the foregoing, it will be appreciated that, although specific  
10 embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

## CLAIMS

1. A method for determining the risk for or presence of a disease associated with altered mitochondrial function in a first subject suspected of having or being at risk for having such a disease, comprising:

comparing a ratio  $r$  for each of a first and a second biological sample containing extramitochondrial DNA and mitochondrial DNA, said first biological sample being obtained from said first subject and said second sample being obtained from a second subject known to be free of a risk or presence of a disease associated with altered mitochondrial function, using the formula:

$$r = x/(x + y)$$

wherein

$x$  is the amount of extramitochondrial DNA in a sample, and

$y$  is the amount of mitochondrial DNA in the sample; and

therefrom determining the risk or presence of the disease.

2. The method of claim 1 wherein the ratio  $r$  is calculated by a method comprising:

contacting a biological sample containing extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in said extramitochondrial DNA and present in said mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of said primer to the extramitochondrial DNA and to the mitochondrial DNA; and

detecting hybridization of the primer to the extramitochondrial DNA and to the mitochondrial DNA, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA.

3. The method of claim 1 wherein the ratio  $r$  is calculated by a method comprising:

contacting a sample containing amplified extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in said amplified extramitochondrial DNA and present in said amplified mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of said primer to the extramitochondrial DNA and to the mitochondrial DNA; and

detecting hybridization of the primer to the extramitochondrial DNA and to the mitochondrial DNA, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA.

4. The method of claim 1 wherein the ratio  $r$  is calculated by a method comprising:

contacting a biological sample containing extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in said extramitochondrial DNA and present in said mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of said primer to the extramitochondrial DNA and to the mitochondrial DNA; and

detecting hybridization and extension of the primer to the extramitochondrial DNA to produce a first product and hybridization and extension of the primer to the mitochondrial DNA to produce a second product distinguishable from said first product, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA.

5. The method of claim 1 wherein the ratio  $r$  is calculated by a method comprising:

contacting a sample containing amplified extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in said amplified extramitochondrial DNA and present in said amplified mitochondrial DNA, under conditions and for a time sufficient to allow

hybridization of said primer to the extramitochondrial DNA and to the mitochondrial DNA:  
and

detecting hybridization and extension of the primer to the extramitochondrial DNA to produce a first product and hybridization and extension of the primer to the mitochondrial DNA to produce a second product distinguishable from said first product, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA.

6. The method of claim 1 wherein the biological sample is treated by heating said sample in water to lyse cells contained in the sample, and then extracting cellular DNA from said lysed cells using an aqueous DNA extraction procedure.

7. The method of claim 6 wherein the sample comprises a crude buffy coat fraction of whole blood.

8. The method of claim 1, further comprising the step of determining the ApoE genotype of the first subject and correlating said genotype with the risk or presence of disease.

9. The method of claim 1 wherein the disease associated with altered mitochondrial function is selected from the group consisting of Alzheimer's Disease, Huntington's Disease, Parkinson's Disease, dystonia, schizophrenia, non-insulin dependent diabetes mellitus, mitochondrial encephalopathy, lactic acidosis, and stroke, myoclonic epilepsy ragged red fiber syndrome, and Leber's hereditary optic neuropathy.

10. A method for quantifying extramitochondrial DNA, comprising:  
contacting a sample containing extramitochondrial DNA with an oligonucleotide primer having a nucleotide sequence complementary to at least a portion of said extramitochondrial DNA under conditions and for a time sufficient to allow hybridization of said primer to the extramitochondrial DNA; and

detecting hybridization of the primer to the extramitochondrial DNA, and therefrom quantifying the extramitochondrial DNA.

11. A method for quantifying extramitochondrial DNA, comprising:

contacting a sample containing extramitochondrial DNA with an oligonucleotide primer having a nucleotide sequence complementary to at least a portion of said extramitochondrial DNA under conditions and for a time sufficient to allow hybridization of said primer to the extramitochondrial DNA; and

detecting hybridization and extension of the primer to the extramitochondrial DNA to produce a product, and therefrom quantifying the extramitochondrial DNA.

12. A method for quantifying extramitochondrial DNA, comprising:

contacting a sample containing amplified extramitochondrial DNA with an oligonucleotide primer having a nucleotide sequence complementary to at least a portion of said extramitochondrial DNA under conditions and for a time sufficient to allow hybridization of said primer to the extramitochondrial DNA; and

detecting hybridization of the primer to the extramitochondrial DNA, therefrom quantifying the extramitochondrial DNA.

13. A method for quantifying extramitochondrial DNA, comprising:

contacting a sample containing amplified extramitochondrial DNA with an oligonucleotide primer having a nucleotide sequence complementary to at least a portion of said extramitochondrial DNA under conditions and for a time sufficient to allow hybridization of said primer to the extramitochondrial DNA; and

detecting hybridization and extension of the primer to the extramitochondrial DNA to produce a product, and therefrom quantifying the extramitochondrial DNA.

14. The method of claim 12 wherein the extramitochondrial DNA is

amplified using a technique selected from the group consisting of polymerase chain reaction, transcriptional amplification systems and self-sustained sequence replication.



15. The method of any one of claims 1-14 wherein a single oligonucleotide primer is used.

16. The method of any one of claims 1-14 wherein a primer extension assay is used.

17. The method of either claim 12 or 14 wherein the step of detecting comprises a technique selected from the group consisting of polymerase chain reaction, primer extension assay, ligase chain reaction, and restriction fragment length polymorphism analysis.

18. The method of any one of claims 1, 10, 11, 12 or 13 wherein the amount of extramitochondrial DNA in the biological sample is quantified by determining the presence in said sample of a nucleotide sequence selected from the group consisting of:

- (a) SEQ ID NO:1,
- (b) a portion of SEQ ID NO:1,
- (c) SEQ ID NO:3,
- (d) a portion of SEQ ID NO:3,
- (e) an extramitochondrial DNA sequence comprising a nucleic acid sequence that (i) corresponds to at least a portion of SEQ ID NO:2 and (ii) contains at least one nucleotide substitution of Figure 4 at a corresponding nucleotide position, and
- (f) an extramitochondrial DNA sequence comprising a nucleic acid sequence that (i) corresponds to at least a portion of SEQ ID NO:2 and (ii) contains at least one nucleotide deletion of Figure 4 at a corresponding nucleotide position.

19. The method of claim 18 wherein the portion of the nucleotide sequence of SEQ ID NO:1 corresponds to a portion of the nucleotide sequence of SEQ ID NO:2 encoding a mitochondrial cytochrome c oxidase.

20. The method of claim 18 wherein the portion of SEQ ID NO:1 corresponds to a portion of a mitochondrial cytochrome c oxidase encoding sequence selected from the group consisting of a portion of a CO1 encoding sequence and a portion of a CO2 encoding sequence.

21. The method of claim 18 wherein the portion of the nucleotide sequence of SEQ ID NO:1 corresponds to a portion of the nucleotide sequence of SEQ ID NO:2 encoding a mitochondrial ATP synthetase subunit.

22. The method of claim 18 wherein the portion of SEQ ID NO:1 corresponds to a portion of a mitochondrial ATP synthetase subunit encoding sequence selected from the group consisting of a portion of a sequence encoding ATP synthetase subunit 6 and a portion of a sequence encoding ATP synthetase subunit 8.

23. The method of claim 18 wherein the nucleotide sequence of SEQ ID NO:1 corresponds to a portion of SEQ ID NO:2 selected from the group consisting of a portion of a sequence encoding ND1, a sequence encoding a portion of ND2 and a sequence encoding a portion of CO3.

24. The method of claim 18 wherein the portion of the nucleotide sequence of SEQ ID NO:3 corresponds to a portion of the nucleotide sequence of SEQ ID NO:2 encoding a mitochondrial ATP synthetase subunit.

25. The method of claim 18 wherein the portion of SEQ ID NO:3 corresponds to a portion of a mitochondrial ATP synthetase subunit encoding sequence selected from the group consisting of a portion of a sequence encoding ATP synthetase subunit 6 and a portion of a sequence encoding ATP synthetase subunit 8.

26. The method of claim 18 wherein the portion of the nucleotide sequence of SEQ ID NO:1 corresponds to a portion of the nucleotide sequence of SEQ ID NO:2 encoding a mitochondrial tRNA.

27. The method of claim 18 wherein the portion of the nucleotide sequence of SEQ ID NO:3 corresponds to a portion of the nucleotide sequence of SEQ ID NO:2 encoding a mitochondrial tRNA.

28. An isolated nucleic acid comprising all or a portion of the nucleotide sequence of SEQ ID NO:1 or a complementary sequence thereto.

29. An isolated nucleic acid comprising all or a portion of a nucleotide sequence of SEQ ID NO:1 or a complementary sequence thereto, wherein the sequence of said isolated nucleic acid differs by at least one nucleotide from the corresponding sequence of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:2 or a complementary sequence thereto.

30. An isolated nucleic acid comprising all or a portion of the nucleotide sequence of SEQ ID NO:3 or a complementary sequence thereto.

31. An isolated nucleic acid comprising all or a portion of a nucleotide sequence of SEQ ID NO:3 or a complementary sequence thereto, wherein the sequence of said isolated nucleic acid differs by at least one nucleotide from the corresponding sequence of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:2 or a complementary sequence thereto.

32. A method for determining the risk or presence of a disease associated with altered mitochondrial function in a subject suspected of having or being at risk for having such a disease, comprising:

quantifying the amount of extramitochondrial DNA and the amount of mitochondrial DNA in a biological sample from said subject, and therefrom determining the risk or presence of the disease.

33. A method for determining the risk or presence of a disease associated with altered mitochondrial function in a first subject suspected of having or being at risk for having such a disease, comprising:

comparing the amount of extramitochondrial DNA and the amount of mitochondrial DNA in a biological sample from said first subject to the amount of extramitochondrial DNA and the amount of mitochondrial DNA in a biological sample from a second subject, and therefrom determining the risk or presence of the disease.

34. A method for determining the risk or presence of a disease associated with altered mitochondrial function in a first subject suspected of having or being at risk for having such a disease, comprising:

quantifying the amount of extramitochondrial DNA and the amount of mitochondrial DNA in a biological sample from said subject and comparing said amount of extramitochondrial DNA and said amount of mitochondrial DNA to the amount of extramitochondrial DNA and the amount of mitochondrial DNA in a biological sample from a second subject known to be free of a risk or presence of a disease associated with altered mitochondrial function, and therefrom determining the risk or presence of the disease.

35. A method of monitoring a telomere comprising:

detecting in a telomeric region the presence of a nucleic acid molecule comprising all or a portion of SEQ ID NO:1 or a complementary portion thereto.

36. The method of claim 35 wherein the detected nucleic acid molecule comprises an exmtDNA sequence.

37. A method of monitoring a telomere comprising:  
detecting in a telomeric region the presence of a nucleic acid molecule comprising all or a portion of SEQ ID NO:3 or a complementary portion thereto.

38. A method of identifying an agent suitable for treating a disease associated with altered mitochondrial function, comprising:

comparing a ratio  $r$  from a sample obtained before contacting a biological source with a candidate agent to the ratio  $r$  from a sample obtained after contacting the biological source with the candidate agent, said ratio  $r$  calculated using the formula:

$$r = x/(x + y)$$

wherein

$x$  is the amount of extramitochondrial DNA in a sample, and

$y$  is the amount of mitochondrial DNA in the sample; and

therefrom determining the suitability of said candidate agent for treating a disease associated with altered mitochondrial function.

39. The method of claim 38 wherein the biological sample comprises a crude buffy coat fraction of whole blood.

40. The method of claim 39 wherein the biological sample is treated by heating said sample in water to lyse cells contained in the sample, and then extracting cellular DNA from said lysed cells using an aqueous DNA extraction procedure.

41. The method of claim 38 wherein the ratio  $r$  is calculated by a method comprising:

contacting a sample containing extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in said extramitochondrial DNA and present in said mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of said primer to the extramitochondrial DNA and to the mitochondrial DNA; and

detecting hybridization of the primer to the extramitochondrial DNA and to the mitochondrial DNA, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA to calculate the ratio  $r$ .

42. The method of claim 38 wherein the ratio  $r$  is calculated by a method comprising:

contacting a sample containing extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in said extramitochondrial DNA and present in said mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of said primer to the extramitochondrial DNA and to the mitochondrial DNA; and

detecting hybridization and extension of the primer to the extramitochondrial DNA to produce a first product and hybridization and extension of the primer to the mitochondrial DNA to produce a second product distinguishable from said first product, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA to calculate the ratio.

43. The method of claim 38 wherein the ratio  $r$  is calculated by a method comprising:

contacting a sample containing amplified extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in said amplified extramitochondrial DNA and present in said amplified mitochondrial DNA, under conditions and for a time sufficient to allow

hybridization of said primer to the extramitochondrial DNA and to the mitochondrial DNA;  
and

detecting hybridization of the primer to the extramitochondrial DNA and to the mitochondrial DNA, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA to calculate the ratio  $r$ .

44. The method of claim 38 wherein the ratio  $r$  is calculated by a method comprising:

contacting a sample containing amplified extramitochondrial DNA and mitochondrial DNA with an oligonucleotide primer having a nucleotide sequence that is complementary to a sequence present in said amplified extramitochondrial DNA and present in said amplified mitochondrial DNA, under conditions and for a time sufficient to allow hybridization of said primer to the extramitochondrial DNA and to the mitochondrial DNA;  
and

detecting hybridization and extension of the primer to the extramitochondrial DNA to produce a first product and hybridization and extension of the primer to the mitochondrial DNA to produce a second product distinguishable from said first product, and therefrom quantifying the extramitochondrial DNA and the mitochondrial DNA to calculate the ratio.

45. The method of claim 38 wherein comparing the ratio  $r$  from a sample obtained before contacting a biological source with a candidate agent to the ratio  $r$  from a sample obtained after contacting the biological source with the candidate agent comprises determination of the presence in said sample of a nucleotide sequence selected from the group consisting of:

- (a) SEQ ID NO:1,
- (b) a portion of SEQ ID NO:1,
- (c) SEQ ID NO:3,
- (d) a portion of SEQ ID NO:3.

(e) an extramitochondrial DNA sequence comprising a nucleic acid sequence that (i) corresponds to at least a portion of SEQ ID NO:2 and (ii) contains at least one nucleotide substitution of Figure 4 at a corresponding nucleotide position, and

(f) an extramitochondrial DNA sequence comprising a nucleic acid sequence that (i) corresponds to at least a portion of SEQ ID NO:2 and (ii) contains at least one nucleotide deletion of Figure 4 at a corresponding nucleotide position.

46. The method of claim 45 wherein the nucleotide sequence of SEQ ID NO:1 or a portion thereof corresponds to a mitochondrial cytochrome c oxidase encoding sequence of SEQ ID NO:2 or a portion thereof.

47. The method of claim 49 wherein the mitochondrial cytochrome c oxidase encoding sequence of SEQ ID NO:2 or a portion thereof is selected from the group consisting of a sequence encoding CO I or a portion thereof and a sequence encoding CO2 or a portion thereof.

48. The method of claim 45 wherein the nucleotide sequence of SEQ ID NO:1 or portion thereof, or the nucleotide sequence of SEQ ID NO:3 or portion thereof corresponds to a mitochondrial ATP synthetase subunit encoding sequence of SEQ ID NO:2 or a portion thereof.

49. The method of claim 48 wherein the mitochondrial ATP synthetase subunit encoding sequence of SEQ ID NO:2 or a portion thereof is selected from the group consisting of a sequence encoding ATP synthetase subunit 6 or a portion thereof and a sequence encoding ATP synthetase subunit 8 or a portion thereof.

50. The method of claim 45 wherein the nucleotide sequence of SEQ ID NO:1 corresponds to a sequence of SEQ ID NO:2 or a portion thereof selected from the group consisting of a sequence encoding a truncated NADH dehydrogenase subunit 1 or a portion



thereof, a sequence encoding NADH dehydrogenase subunit 2 or a portion thereof and a sequence encoding truncated CO3 or a portion thereof.

51. The method of claim 38 wherein the disease associated with altered mitochondrial function is selected from the group consisting of Alzheimer's Disease, Huntington's Disease, Parkinson's Disease, dystonia, schizophrenia, non-insulin dependent diabetes mellitus, mitochondrial encephalopathy, lactic acidosis, and stroke, myoclonic epilepsy ragged red fiber syndrome, and Leber's hereditary optic neuropathy.

52. A method of identifying an agent suitable for treating a subject suspected of being at risk for having a disease associated with altered mitochondrial function, comprising:

determining the apolipoprotein E genotype of the subject;

comparing a ratio  $r$  in a biological sample obtained from the subject before contacting said sample with a candidate agent to the ratio  $r$  in a biological sample obtained from the subject after contacting said sample with a candidate agent, said ratio  $r$  calculated using the formula:

$$r = x / (x + y)$$

wherein

$x$  is the amount of extramitochondrial DNA in the sample, and

$y$  is the amount of mitochondrial DNA in the sample; and

therefrom determining the suitability of said candidate agent for treating the disease associated with altered mitochondrial function.

53. The method of claim 52 wherein the disease associated with altered mitochondrial function is Alzheimer's disease.

54. A method of correlating a ratio  $r$  with the suitability of an agent for treating Alzheimer's disease in a subject, comprising:

determining a ratio  $r$  in a biological sample obtained from the subject. said ratio  $r$  calculated using the formula:

$$r = x / (x + y)$$

wherein

$x$  is the amount of extramitochondrial DNA in the sample, and

$y$  is the amount of mitochondrial DNA in the sample;

contacting said subject with a candidate agent and evaluating the subject for alterations in the AD disease state, and therefrom correlating the suitability of the agent for treating AD in the subject with  $r$ .

55. The method of claim 54, further comprising determining the apolipoprotein E genotype of the subject, and therefrom correlating the suitability of the agent for treating AD in the subject with  $r$  and with the apolipoprotein E genotype.

1/42

cactgtaaag	ctaaccagc	attaaccttt	taagttaaag	actaagagaa	tcattatctc	60
tttacagtga	aatgccacag	ctaaatacca	ctgtatgacc	tgctatcatc	acccaatac	120
tcctcacgtt	atttctcatc	acccaactaa	aaatactaaa	cacacactgc	catctgcca	180
cctcacaaa	atttattaaa	ataaaaaact	acagtaagcc	ctgagaacca	aatgaacga	240
aaatttattc	gcttcattca	ttacccttac	agtactaggc	ctaccgcca	cagtaccaat	300
catcctattt	cccccttac	tggtccaac	ctccaaatac	ctcatcaaca	accgactaat	360
caccactcaa	caatgactac	ttcaactcac	cttaaaacaa	ataataacga	tacataacat	420
taagggacga	acctggtccc	ttatacta	ttccctgatt	atttttattg	ccacaactaa	480
tctcctcgga	ctcttgcccc	actcatttac	accaatcac			519

*Fig. 1*

*Fig. 2A*

gatcacagg	ctatcaccc	attaaccact	cacgggagct	ctccatgcat	ttggattttt	60
cgtctggggg	gtatgcacgc	gatagcattg	cgagacgctg	gagccggagc	accctatgtc	120
gcagtatctg	tctttgattc	ctgcctcatc	ctattattta	tcgcacctac	gttcaatatt	180
acaggcgaac	atacttacta	aagtgtgtta	attaattaat	gcttgttagga	cataataata	240
acaattgaat	gtctgcacag	ccactttcca	cacagacatc	ataacaaaaa	atttccacca	300
aacccccctt	cccccgcttc	tggccacagc	acttaaacac	atctctgcca	aacccccaaa	360
acaaagaacc	ctaaccaccg	cctaaccaga	tttcaaattt	tatcttttgg	cggtatgcac	420
ttttaacagt	caccccccaa	ctaacacatt	attttcccct	cccactccca	tactactaat	480
ctcatcaata	caacccccgc	ccatcctacc	cagcacacac	acaccgctgc	taaccccata	540
ccccgaacca	accaaaccct	aaagacaccc	cccacagttt	atgtagctta	cctcctcaaa	600
gcaatacact	gaaaaatgtt	agacgggctc	acatcacccc	ataaacaaat	aggtttggct	660
ctagcctttc	tattagctct	tagtaagatt	acacatgcaa	gcatccccgt	tccagtgaat	720
tcaccctcta	aatcaccacg	atcaaaaagg	acaagcatca	agcacgcagc	aatgcagctc	780
aaaacgctta	gcctagccac	acccccacgg	gaaacagcag	tgattaacct	ttagcaataa	840
acgaaagttt	aactaagcta	tactaacccc	aggttggtgc	aatttcgtgc	cagccaccgc	900
ggtcacacga	tttaacccaag	tcaatagaag	ccggcgtaaa	gagtgtttta	gatcaccccc	960
tccccaataa	agctaaaact	cacctgagtt	gtaaaaaact	ccagttgaca	caaaatagac	1020
tacgaaagtg	gctttaacat	atctgaacac	acaatagcta	agacccaaac	tgggattaga	1080
tacccccacta	tgcttagccc	taaaacctcaa	cagttaaatc	aacaaaaactg	ctcgcagaa	1140
cactacgagc	cacagcttaa	aactcaaaagg	acctggcggg	gcttcataatc	cctctagagg	1200
agcctgttct	gtaatcgata	aacccccgatc	aacctcacca	cctcttgctc	agcctatata	1260
ccgccatctt	cagcaaacc	tgatgaaggc	tacaaagtaa	gcgcaagtac	ccacgtaaa	1320
acgttaggtc	aaggtgtagc	ccatgaggtg	gcaagaaatg	ggctacattt	tctaccccc	1380
aaaactacga	tagcccttat	gaaacttaag	ggtcgaagg	ggatttagca	gtaaactaag	1440
agtagagtgc	ttagttgaac	agggccctga	agcgctgaca	caccgcccgt	caccctcctc	1500
aagtatactt	caaaggacat	ttaactaaaa	ccccctacgca	tttatataga	ggagacaagt	1560
cgtaacattg	taagtgtact	ggaaagtgc	cttggacgaa	ccagagtgtg	gcttaacaca	1620
aagcacccaa	cttacactta	ggagatttca	acttaacttg	accgctctga	gctaaacct	1680
gccccaaacc	cactccacct	tactaccaga	caaccttagc	caaaccattt	acccaaataa	1740
agtataggcg	atagaaattg	aaacctggcg	caatagatat	agtaccgcaa	gggaaagatg	1800
aaaaattata	accaagcata	atatagcaag	gactaacccc	tataccttct	gcataatgaa	1860
ttaactagaa	ataactttgc	aaggagagcc	aaagctaaga	cccccgaaac	cagacgagct	1920
acctaagaac	agctaaaaga	gcacaccggt	ctatgtagca	aaatagtggg	aagatttata	1980
ggtagaggcg	acaaacctac	cgagcctggt	gatagctggt	tgtccaagat	agaatcttag	2040
ttcaacttta	aatttgccca	cagaaccttc	taaatcccct	tgtaaattta	actgttagtc	2100
caaagaggaa	cagctctttg	gacactagga	aaaaaccttg	tagagagagt	aaaaaattta	2160
acacccatag	taggcctaaa	agcagccacc	aattaagaaa	gcgttcaagc	tcaacaccca	2220
ctacctaaaa	aatcccaaac	atataactga	actcctcaca	cccaattgga	ccaatctatc	2280
accttataga	agaactaatg	ttagtataag	taacatgaaa	acattctcct	ccgcataagc	2340
ctgcgtcaga	ttaaaacact	gaactgacaa	ttacagccc	aatatctaca	atcaaccaac	2400
aagtcattat	taccctcact	gtcaacccaa	cacaggcatg	ctcataagga	aaggttaaaa	2460
aaagtaaaa	gaactcggca	aatcttacc	cgctgtttta	ccaaaaacat	cacctctagc	2520
atcaccagta	ttagaggcac	cgctgccc	gtgacacatg	tttaacggcc	gcggtaccct	2580
aaccgtgcaa	aggtagcata	atcacttggt	ccttaaatag	ggacctgtat	gaatggctcc	2640
acgagggttc	agctgtctct	tacttttaac	cagtgaat	gacctgccc	tgaagaggcg	2700
ggcataaac	agcaagacga	gaagacccta	tggagcttta	atttattaat	gcaaacagta	2760
cctaacaaac	ccacaggtcc	taaactacca	aacctgcatt	aaaaatttcg	gttggggcga	2820
cctcggagca	gaacccaacc	tccgagcagt	acatgctaag	acttcaccag	tcaaagcgaa	2880
ctactatact	caattgatcc	aataacttga	ccaacggaac	aagttaccct	agggataaca	2940
gcgcaatcct	attctagagt	ccatatcaac	aatagggttt	acgacctcga	tggttgatca	3000
ggacatccc	atggtgcagc	cgctattaaa	ggttcggttg	ttcaacgatt	aaagtcctac	3060
gtgatctgag	ttcagaccgg	agtaatccag	gtcgggtttc	atctaccttc	aaattcctcc	3120
ctgtacgaaa	ggacaagaga	aataaggcct	acttcacaaa	gcgccttccc	ccgtaaatga	3180
tatcatctca	acttagtatt	ataccacac	ccaccaaga	acagggttg	tttaagatggc	3240
agagcccgg	aatcgcataa	aacttaaaac	tttacagtca	gaggttcaat	tcctcttctt	3300
aaacaatac	ccatggccaa	cctcctactc	ctcattgtac	ccattcta	cgcaatggca	3360
ttcctaattgc	ttaccgaacg	aaaaattcta	ggctatatac	aactacgcaa	agggcccaac	3420
gttgtagggc	cctacgggct	actacaacc	ttcgtgacg	ccataaaact	cttcacaaa	3480

*Fig. 2B*

3/42

gagccccctaa	aaccgcgcac	atctaccatc	accctctaca	tcaccgcccc	gaccttagct	3540
ctcaccatcg	ctcttctact	atgaaccccc	ctccccatac	ccaacccccct	ggtcaacctc	3600
aacctaggcc	tcctatttat	tctagccacc	tctagccctag	ccgtttactc	aatcctctga	3660
tcagggtgag	catcaaactc	aaactacgcc	ctgatcggcg	cactgcgagc	agtagcccaa	3720
acaatctcat	atgaagtcac	cctagccatc	attctactat	caacattact	aataagtggc	3780
tcctttaacc	tctccaccct	tatcacaaca	caagaacacc	tctgattact	cctgccatca	3840
tgacccttgg	ccataatatg	atttatctcc	acactagcag	agaccaaccg	aacccccctc	3900
gaccttgccg	aaggggagtc	cgaactagtc	tcaggcttca	acatcgaaata	cgccgcagggc	3960
cccttcgcgc	tattcttcat	agccgaatac	acaaacatta	ttataataaa	cacccctacc	4020
actacaatct	tcctaggaac	aacatatgac	gcactctccc	ctgaactcta	cacaacatat	4080
tttgtcacca	agaccctact	tctaacctcc	ctgttcttat	gaattcgaac	agcatacccc	4140
cgattccgct	acgaccaact	catacacctc	ctatgaaaaa	acttcctacc	actcacctta	4200
gcattactta	tatgatatgt	ctccataccc	attacaatct	ccagcattcc	ccctcaaacc	4260
taagaaatat	gtctgataaa	agagttactt	tgatagagta	aataatagga	gcttaaaccc	4320
ccttattttct	aggactatga	gaatcgaacc	catccctgag	aatccaaaat	tctccgtgcc	4380
acctatcaca	ccccatccta	aagtaaggtc	agctaaataa	gctatcgggc	ccataccccg	4440
aaaatgttgg	ttataccctt	cccgtactaa	ttaatccctt	ggcccaaccc	gtcatctact	4500
ctaccatctt	tgcaggcaca	ctcatcacag	cgctaagctc	gcactgattt	tttacctgag	4560
taggcctaga	aataaacatg	ctagctttta	ttccagttct	aaccaaaaaa	ataaaccttc	4620
gttccacaga	agctgccatc	aagtatttcc	tcacgcaagc	aaccgcatcc	ataatccttc	4680
taatagctat	cctcttcaac	aatatactct	ccggacaatg	aaccataacc	aatactacca	4740
atcaatactc	atcattaata	atcataatag	ctatagcaat	aaaactagga	atagccccct	4800
ttcactttctg	agtcccgag	gttaccctct	gcaccctctt	gacatccggc	ctgcttcttc	4860
tcacatgaca	aaaactagcc	cccattctca	tcataatcca	aatctctccc	tcactaaacg	4920
taagccttct	cctcactctc	tcaatcttat	ccatcatagc	aggcagttga	ggtggattaa	4980
accagacca	gctacgcaaa	atcttagcat	actcctcaat	taccacata	ggatgaataa	5040
tagcagttct	accgtacaac	cctaacataa	ccattcttaa	tttaactatt	tatattatcc	5100
taactactac	cgcattccta	ctactcaact	taaactccag	caccacgacc	ctactactat	5160
ctcgcacctg	aaacaagcta	acatgactaa	cacccttaat	tccatccacc	ctcctctccc	5220
taggaggcct	gccccgccta	accggctttt	tgcccaaatg	ggccattatc	gaagaattca	5280
caaaaaacaa	tagcctcatc	atccccacca	tcatagccac	catcacccctc	cttaacctct	5340
acttctacct	acgcctaata	tactccacct	caatcacact	actccccata	tctaacaacg	5400
taaaaataaa	atgacagttt	gaacatacaa	aaccaccccc	attcctcccc	acactcatcg	5460
cccttaccac	gctactccta	cctatctccc	cttttatact	aataatctta	tagaaattta	5520
ggttaaatac	agaccaagag	ccttcaaagc	cctcagtaag	ttgcaatact	taatttctgt	5580
aacagctaag	gactgcaaaa	ccccactctg	catcaactga	acgcaaatca	gccactttaa	5640
ttaagctaag	cccttactag	accaatggga	cttaaaccca	caaacactta	ggttaacagct	5700
aagcaccccta	atcaactggc	ttcaatctac	ttctcccgcg	gccgggaaaa	aaggcgaggag	5760
aagccccggc	aggtttgaag	ctgcttcttc	gaatttgcaa	ttcaatatga	aaatcacctc	5820
ggagctggta	aaaagaggcc	taaccctgtt	cttttagattt	acagtccaat	gcttcactca	5880
gccatttttac	ctcaccccca	ctgatgttcg	ccgaccgttg	actatttctt	acaaaccaca	5940
aagacattgg	aacactatac	ctattattcg	gcgcatgagc	tgagtccta	ggcacagctc	6000
taagcctcct	tattcgagcc	gagctgggcc	agccaggcaa	ccttctaggt	aacgaccaca	6060
tctacaacgt	tatcgtcaca	gcccattgcat	ttgtaataat	cttcttcata	gtaataacca	6120
tcataatcgg	aggctttggc	aactgactag	ttcccctaata	aatcggtgcc	cccgatatgg	6180
cgtttccccg	cataaacaac	ataagcttct	gactcttacc	tccctctctc	ctactcctgc	6240
tcgcatctgc	tatagtggag	gccggagcag	gaacagggtg	aacagtctac	cctcccttag	6300
cagggaaacta	ctcccaccct	ggagcctccg	tagacctaac	catcttctcc	ttacacctag	6360
caggtgtctc	ctctatctta	ggggccatca	atttcatcac	aacaattatc	aatataaaaac	6420
cccctgccat	aacccaatac	caaacgcccc	tcttcgtctg	atccgtccta	atcacagcag	6480
tcctacttct	cctatctctc	ccagtcctag	ctgttgccat	cactatacta	ctaacagacc	6540
gcaacctcaa	caccaccttc	ttcggacccg	ccggaggagg	agaccccatc	ctataccaac	6600
acctattctg	atTTTTcggt	caccctgaag	tttatattct	tatcctacca	ggcttcggaa	6660
taatctccca	tattgtaact	tactactccg	gaaaaaaaga	accatttgga	tacataggta	6720
tggctctgagc	tatgatatca	attggcttcc	taggggttat	cgtgtgagca	caccatatat	6780
ttacagtagg	aatagacgta	gacacacgag	catatttcac	ctccgctacc	ataatcatcg	6840
ctatccccac	cggcgtcaaa	gtatttagct	gactcgccac	actccacgga	agcaatatga	6900
aatgatctgc	tgcagtgtct	tgagccctag	gattcatctt	tcttttcacc	gtaggtggcc	6960

*Fig. 2C*

tgactggcat	tgtattagca	aactcatcac	tagacatcgt	actacacgac	acgtactacg	7020
ttgtagccca	cttccactat	gtcctatcaa	taggagctgt	atttgccatc	ataggaggct	7080
tcattcactg	atttccccta	ttctcaggct	acaccctaga	ccaaacctac	gccaaaatcc	7140
atttcactat	catattcatc	ggcgtaaate	taactttctt	cccacaacac	tttctcggcc	7200
tatccggaat	gccccgacgt	tactcggact	accccgatgc	atacaccaca	tgaacatcc	7260
tatcatctgt	aggctcattc	atctctctaa	cagcagtaat	attaataatt	ttcatgattt	7320
gagaagccct	cgcttcgaag	cgaaaagtcc	taatagtaga	agaaccctcc	ataaacctgg	7380
agtgactata	tggatgcccc	ccaccctacc	acacattcga	agaaccctga	tacataaaat	7440
ctagacaaaa	aaggaaaggaa	tcgaaccccc	caaagctggg	ttcaagccaa	ccccatggcc	7500
tccatgactt	tttcaaaaag	gtattagaaa	aaccatttca	taactttgtc	aaagttaaat	7560
tataggctaa	atcctatata	tcttaatggc	acatgcagcg	caagtaggtc	tacaagacgc	7620
tacttcccct	atcatagaag	agcttatcac	ctttcatgat	cacgcccctca	taatcatttt	7680
ccttatctgc	ttcctagtcc	tgtatgcctt	tttccctaaca	ctcacaacaa	aactaactaa	7740
tactaacaatc	tcagacgctc	aggaataga	aaccgtctga	actatcctgc	ccgccatcat	7800
cctagtccct	atcgccctcc	catccctacg	catcctttac	ataacagacg	aggtcaacga	7860
tccctccctt	accatcaaat	caattggcca	ccaatggtac	tgaacctacg	agtaaccgga	7920
ctacggcgga	ctaattcttca	actcctacat	acttccccca	ttattcctag	aaccaggcga	7980
cctgacgctc	cttgacgttg	acaatcgagt	agtactcccg	attgaagccc	ccattcgtat	8040
aataattaca	tcacaagacg	tcttgacttc	atgagctgtc	cccacattag	gcttaaaaac	8100
agatgcaatt	cccggacgtc	taaaccaaac	cactttcacc	gctacacgac	cgggggtata	8160
ctacgggtcaa	tgctctgaaa	tctgtggagc	aaaccacagt	ttcatgcccc	tcgtcctaga	8220
attaattccc	ctaaaaatct	ttgaaatagg	gcccgtatit	accctatagc	accccctcta	8280
ccccctctag	agcccactgt	aaagctaact	tagcattaac	cttttaagtt	aaagattaag	8340
agaaccaaca	cctctttaca	gtgaaatgcc	ccaactaaat	actaccgtat	ggcccaccat	8400
aattaccccc	atactcctta	cactattcct	catcacccaa	ctaaaaatat	taaacacaaa	8460
ctaccaccta	cctccctcac	caaagcccat	aaaaataaaa	aattataaca	aaccctgaga	8520
acaaaatga	acgaaaatct	gttcgcttca	ttcattgccc	ccacaatcct	aggcctacc	8580
gccgcagtac	tgatcattct	atctccccct	ctattgatcc	ccacctccaa	atatctcatc	8640
aacaaccgac	taatcaccac	ccaacaatga	ctaattcaaa	taacctcaaa	acaaatgata	8700
accatacaca	acactaaagg	tctctttatac	tctctttatac	tagtatcctt	aatcatTTTT	8760
attgccacaa	ctaacctcct	cggactcctg	cctcactcat	ttacaccaac	cacccaacta	8820
tctataaacc	tagccatggc	catccccctt	tgagcgggca	cagtgattat	aggctttcgc	8880
tctaagatta	aaaatgcctt	agcccacttc	ttaccacaag	gcacacctac	accccttatc	8940
cccatactag	ttattatcga	aaccatcagc	ctactcattc	aaccaatagc	cctggccgta	9000
cgcctaaccg	ctaacattac	tgcaggccac	ctactcatgc	acctaattgg	aagcggcacc	9060
ctagcaatat	caaccattaa	ccttcccttc	acacttatca	tcttcacaat	tctaattcta	9120
ctgacttatcc	tagaaatcgc	tgctgcctta	atccaagcct	acgttttcac	acttctagta	9180
agcctctacc	tgacgacaaa	cacataatga	cccaccaatc	acatgcctat	catatagtaa	9240
aaccagcccc	atgaccccta	acagggggcc	tctcagccct	cctaattgacc	tccggcctag	9300
ccatgtgatt	tcacttccac	tccataacgc	tctcataact	aggcctacta	accaacacac	9360
taaccatata	ccaatgatgg	cgcgatgtaa	cacgagaaaag	cacataccaa	ggccaccaca	9420
caccacctgt	ccaaaaaggc	cttcgatacg	ggataatcct	atttattacc	tcagaagttt	9480
ttttcttcgc	aggatttttc	tgagcctttt	accactccag	cctagcccct	accccccaat	9540
taggagggca	ctggccccc	acaggcattca	ccccgctaaa	tcccctagaa	gtcccactcc	9600
taaaacacatc	cgtattactc	gcatcaggag	tatcaatcac	ctgagctcac	catagtctaa	9660
tagaaaacaa	ccgaaaccaa	ataattcaag	cactgcttat	tacaatttta	ctgggtctct	9720
attttaccct	cctacaagcc	tcagagtact	tcgagtctcc	cttcaccatt	tccgacggca	9780
tctacggctc	aacatttttt	gtagccacag	gcttccacgg	acttcacgtc	attattggct	9840
caactttcct	cactatctgc	ttcatccgcc	aactaatatt	tcactttaca	tccaaacatc	9900
actttggctt	cgaagccgcc	gcctgatact	ggcattttgt	agatgtgggt	tgactatttc	9960
tgtatgtctc	catctattga	tgaggggtctt	actcttttag	tataaatagt	accgttaact	10020
tccaattaac	tagttttgac	aacattcaaa	aaagagtaat	aaacttcgcc	ttaattttaa	10080
taatcaaac	cctcctagcc	ttactactaa	taattattac	attttgacta	ccacaactca	10140
acggctacat	agaaaaatcc	accccttacg	agtgcggctt	cgaccctata	tccccgccc	10200
gcgtcccttt	ctccataaaa	ttcttcttag	tagctattac	cttcttatta	ttgatctag	10260
aaattgccct	ccttttaccc	ctaccatgag	ccctacaac	aactaacctg	ccactaatag	10320
ttatgtcatc	cctcttatta	atcatcatcc	tagccctaag	tctggcctat	gagtgactac	10380
aaaaaggatt	agactgaacc	gaattggtat	atagtttaaa	caaaacgaat	gatttcgact	10440

*Fig. 2D*

cattaaatta	tgataatcat	atttaccaaa	tgcccctcat	ttacataaat	attatactag	10500
catttaccat	ctcacttcta	ggaatactag	tatatcgctc	acacctcata	tcctccctac	10560
tatgcctaga	aggaataata	ctatcgctgt	tcattatagc	tactctcata	accctcaaca	10620
cccactccct	cttagccaat	attgtgccta	ttgccatact	agcttttgcc	gcctgcgaag	10680
cagcgggtggg	cctagcccta	ctagtctcaa	tctccaacac	atatggccta	gactacgtac	10740
ataacctaata	cctactccaa	tgctaaaact	aatcgtecca	acaattatat	tactaccact	10800
gacatgactt	tccaaaaaac	acataatttg	aatcaacaca	accaccacaca	gcctaattat	10860
tagcatcatc	cctctactat	tttttaacca	aatcaacaac	aacctattta	gctgttcccc	10920
aaccttttcc	tccgaccccc	taacaacccc	cctcctaata	ctaactacct	gactcctacc	10980
cctcacaaac	atggcaagcc	aacgccaactt	atccagtgaa	ccactatcac	gaaaaaaaact	11040
ctacctctct	atactaactt	ccctacaaaat	ctccttaatt	ataacattca	cagccacaga	11100
actaatcata	ttttatatct	tcttcgaaac	cacacttatc	cccaccttgg	ctatcatcac	11160
ccgatgaggg	aaccagccag	aacgcctgaa	cgcaggcaca	tacttccctat	tctacaccct	11220
agtaggctcc	cttcccctac	tcatcgcaact	aatttacact	cacaacaccc	tagggtcact	11280
aaacatttcta	ctactcactc	tcaactgccc	agaactatca	aactcctgag	ccaataactt	11340
aatatgacta	gcttacacaa	tagcttttat	agtaaagata	cctctttacg	gactccactt	11400
atgactccct	aaagccccatg	tcgaagcccc	catcgctggg	tcaatagtag	ttggccgagt	11460
actcttaaaa	ctaggcggt	atggtataat	acgcccacaca	ctcatttcta	accccctgac	11520
aaaacacata	gcctacccct	tccctgtact	atccctatga	ggcataatta	taacaagctc	11580
catctgccta	cgacaaacag	acctaataatc	gctcattgca	tactcttcaa	tcagccacat	11640
agccctcgta	gtaacagcca	ttctcatcca	aaccccctga	agcttcaccg	gcgcagtcac	11700
tctcataatc	gcccacgggc	ttacatcctc	attactattc	tgccatgcaa	actcaaacta	11760
cgaacgcact	cacagtcgca	tcataatcct	ctctcaagga	cttcaaactc	tactcccact	11820
aatagctttt	tgatgacttc	tagcaagcct	cgtaaacctc	gccttaccac	ccactattaa	11880
ctactggga	gaactctctg	tgctagtaac	cacgttctcc	tgatcaaata	tcactctcct	11940
acttacagga	ctcaacatac	tagtcacagc	cctatactcc	ctctacatat	ttaccacaac	12000
acaatggggc	tcactcacc	accacattaa	caacataaaa	ccctcattca	cacgagaaaa	12060
caccctcatg	ttcatacacc	tatcccccat	tctctccta	tccctcaacc	ccgacatcat	12120
taccgggttt	tcctcttgta	aatatagttt	aaccataaaa	tcagattgtg	aatctgacaa	12180
cagaggctta	cgacccctta	tttaccgaga	aagctcacia	gaactgctaa	ctcatgcccc	12240
catgtctttt	aacatggctt	tctcaacttt	taaaggataa	cagctatcca	ttggtcttag	12300
gccccaaaaa	ttttgggtgca	actccaaata	aaagtaataa	ccatgcacac	tactataacc	12360
accctaacc	tgacttccct	aattcccccc	atccttacca	ccctcgtaa	ccctaacaaa	12420
aaaaactcat	acccccatta	tgtaaaatcc	attgtcgcat	ccacctttat	tatcagtcct	12480
ttccccacaa	caatattcat	gtgcctagac	caagaagtta	ttatctcgaa	ctgacactga	12540
gccacaaccc	aaacaaccca	gctctcccta	agcttcaaac	tagactactt	ctccataata	12600
ttcatccctg	tagcattgtt	cgttacatgg	tccatcatag	aattctcact	gtgatatata	12660
aactcagacc	caaacattaa	tcagttcttc	aaatatctac	tcattcttct	aattaccata	12720
ctaactcttag	ttaccgctaa	caacctattc	caactgttca	tcggctgaga	gggcgtagga	12780
attatatcct	tcttgctcat	cagttgatga	tacgcccag	cagatgccaa	cacagcagcc	12840
attcaagcaa	tcctatacaa	ccgtatcggc	gatatcgggt	tcatectcgc	cttagcatga	12900
tttatcctac	actccaactc	atgagaccca	caacaataag	cccttctaaa	cgctaatacca	12960
agcctcacc	cactactagg	cctcctccta	gcagcagcag	gcaaatcagc	ccaattaggt	13020
ctccaccct	gactccctc	agccatagaa	ggcccaccc	cagttctcagc	cctactccac	13080
tcaagcacta	tagttgtagc	aggaatcttc	ttactcatcc	gcttccaccc	cttagcagaa	13140
aatagcccac	taatccaaac	tctaactacta	tgcttagggc	ctatcaccac	tctgttcgca	13200
gcagtctgcg	cccttacaca	aaatgacatc	aaaaaaatcg	tagccttctc	cacttcaagt	13260
caactaggac	tcataatagt	tacaatcggc	atcaaccaac	cacacctagc	attcctgcac	13320
atctgtaccc	acgccttctt	caaagccata	ctatttatgt	gctccgggtc	catcatccac	13380
aaccttaaca	atgaacaaga	tattcgaaaa	ataggaggac	tactcaaaac	catacctctc	13440
acttcaacct	ccctcaccat	tggcagccta	gcattagcag	gaataccttt	cctcacaggt	13500
ttctactcca	aagaccacat	catcgaaacc	gcaaacatat	catacacaaa	cgcctgagcc	13560
ctctctatta	ctctcatcgc	tacctccctg	acaagcgct	atagcactcg	aaataattctt	13620
ctcaccctaa	caggtcaacc	tcgcttcccc	acccttacta	acattaacga	aaataacccc	13680
accctactaa	accccatata	acgcctggca	gccggaagcc	tattcgagg	atttctcatt	13740
actaacaaca	tttcccccg	atcccccttc	caaacaacaa	tccccctcta	cctaaaactc	13800
acagccctcg	ctgtcacttt	cctaggactt	ctaacagccc	tagacctcaa	ctacctaac	13860
aacaaactta	aaataaaatc	cccactatgc	acattttatt	tctccaacat	actcggattc	13920

6/42

taccctagca	tcacacaccg	cacaatcccc	tatctaggcc	ttcttacgag	ccaaaacctg	13980
cccctactcc	tcctagacct	aacctgacta	gaaaagctat	tacctaaaac	aatttcacag	14040
caccaaatct	ccacctccat	catcacctca	acccaaaaag	gcataattaa	actttacttc	14100
ctctctttct	tcttcccact	catcctaacc	ctactcctaa	tcacataaacc	tattcccccg	14160
agcaatctca	attacaatat	atacaccaac	aaacaatgtt	caaccagtaa	ctactactaa	14220
tcaacgcccc	taatcataca	aagccccgcg	accaatagga	tcctcccgaa	tcaaccctga	14280
cccctctcct	tcataaatta	ttcagcttcc	tacactatta	aagtttacca	caaccaccac	14340
cccctacatac	tctttcaccc	acagcaccaa	tcctacctcc	atcgctaacc	ccactaaaac	14400
actcaccaag	acctcaaccc	ctgaccccca	tgcctcagga	tactcctcaa	tagccatcgc	14460
tgtagtatat	ccaaagacaa	ccatcattcc	ccctaaataa	attaaaaaaa	ctattaaaac	14520
catataacct	cccccaaaat	tcagaataat	aacacaccgc	accacaccgc	taacaatcaa	14580
tactaaaccc	ccataaatag	gagaaggctt	agaagaaaaa	cccacaaacc	ccattactaa	14640
accacacatc	aacagaaaaca	aagcatacat	cattattctc	gcacggacta	caaccacgac	14700
caatgatatg	aaaaaccatc	gttggtatttc	aactacaaga	acaccaatga	ccccaatacg	14760
caaaattaac	cccctaataa	aatttaattaa	ccactcattc	atcgacctcc	ccacccccatc	14820
caacatctcc	gcatgatgaa	acttcggctc	actccttggc	gcctgcctga	tcctccaaat	14880
caccacagga	ctatttcctag	ccatgcacta	ctcaccagac	gcctcaaccg	ccttttcac	14940
aatcgcccac	atcactcgag	acgtaaaatta	tggctgaatc	atccgctacc	ttcacgcca	15000
tggcgctca	atattcttta	tctgcctctt	cctacacatc	ggcgagggcc	tatattacgg	15060
atcatttctc	tactcagaaa	cctgaaacat	cggcattatc	ctcctgcttg	caactatagc	15120
aacagccttc	ataggctatg	tcctcccgctg	aggccaaata	tcattctgag	gggccacagt	15180
aattacaaac	ttactatccg	ccatcccata	cattgggaca	gacctagttc	aatgaatctg	15240
aggaggctac	tcagtagaca	gtcccaccct	cacacgattc	tttacccttc	acttcacctt	15300
gcccttcatt	attgcagccc	tagcaaacact	ccacctccta	ttcttgcacg	aaacgggatc	15360
aaacaacccc	ctaggaatca	cctcccattc	cgataaaatc	accttcacc	cttactacac	15420
aatcaaagac	gccctcggct	tacttctctt	ccttctctcc	ttaatgacat	taacactatt	15480
ctcacagac	ctcctaggcg	accagacaaa	ttatacccta	gccaaccctc	taaacacccc	15540
tccccacatc	aagccccgaat	gatatttcct	attcgcttac	acaattctcc	gatccgtccc	15600
taacaaacta	ggaggcgctc	ttgccctatt	actatccatc	ctcatcctag	caataatccc	15660
catcctccat	atatccaaac	aacaagcat	aatatttcgc	ccactaagcc	aatcacttta	15720
ttgactccta	gccgcagacc	tcctcattct	aacctgaatc	ggaggacaac	cagtaagcta	15780
cccttttacc	atcattggac	aagtagcatc	cgtactatac	ttcacacaac	tcctaatcct	15840
aataccaact	atctcccctaa	ttgaaaacaa	aataactcaa	tgggcctgtc	cttgtagtat	15900
aaactaatac	accagtcttg	ttaaaccggag	atgaaaacct	ttttccaagg	acaaatcaga	15960
gaaaaagctt	ttaactccac	cattagcacc	caaagctaag	attctaattt	aaactattct	16020
ctgttctttc	atggggaagc	agatttgggt	accacccaag	tattgactca	cccatacaac	16080
accgctatgt	atttcgtaca	ttactgccag	ccaccatgaa	tattgtacgg	taccataaat	16140
acttgaccac	ctgtagtaca	taaaaaccca	atccacatca	aaacccccctc	cccatagctta	16200
caagcaagta	cagcaatcaa	ccctcaacta	tcacacatca	actgcaactc	caaagccacc	16260
cctcacccac	taggatacca	acaaacctac	ccacccttaa	cagtacatag	tacataaagc	16320
catttaccgt	acatagcaca	ttacagtcaa	atcccttctc	gtccccatgg	atgaccccc	16380
tcagataggg	gtcccttgac	caccatctct	cgtgaaatca	atatcccgcga	caagagtgt	16440
actctcctcg	ctccggggcc	ataacacttg	ggggtagcta	aagtgaactg	tatccgacat	16500
ctggttccta	cttcagggtc	ataaagccta	aatagccac	acgttccccct	taaataagac	16560
atcacgatg						16569

*Fig. 2E*



*Fig. 3A*

ataggcattc	caactcatcc	gcttgctgac	gacctccacg	tgatttcaac	aatgatttca	60
aataatttcac	tttttaagtc	agtgtgactt	aagtatgaaa	ttgcctctcc	ctaaagctcc	120
cctaaggcct	aaacagtcgt	cattaccata	gctgtgacag	ggagactgtt	gaatttataa	180
tctattggcc	attcacagca	tagcgtataa	accctagctc	atgatttctt	tgcaatagaa	240
gtgtactttt	tcatacacatt	cccttcacaa	cttactcacc	agacagactt	tgagctctcc	300
tcctggctta	gcctggatcg	tttgaaatgg	tcataccatcc	tttggcccca	atacctaaac	360
taaggtctat	gaacaataag	atgattttac	ttcagtggga	cttttttggt	taatatatta	420
gatttgacct	tcagcaaggt	caaagggagt	ccgaactagt	ctcaggcttc	aacatcgaat	480
acgccgcagg	ncccttcgcc	ctattcttca	tagccgaata	cacaaacatt	attataataa	540
acacctcac	cactacaatc	ttcctaggaa	caacatataa	cgcactctcc	cctgaactct	600
acacaacata	ttttgtcacc	aagaccctac	ttctgacctc	cctgttctta	tgaattcgaa	660
cagcataccc	ccgattccgc	tacgaccaac	tcatacacct	cctatgaaaa	aacttcctac	720
cactcacctt	agcattactt	ataatgatag	tctccatacc	cattacaatc	tccagcattc	780
cccctcaaac	ctaagaaata	tgtctgataa	aagagttact	ttgatagagt	aaataatagg	840
agtttaaatc	cccttatttc	taggactatg	agaatcgaac	ccatccctga	gaatccaaaa	900
ttctcgggtc	caactatcac	accccatctt	aaagtaagggt	cagctaaata	agctatcggg	960
cccatacccc	gaaaatgttg	gttatatcct	tcccgtacta	attaatcccc	tgccccaacc	1020
cgtcatttac	tctaccatct	ttgcaggcac	actcatcaca	gcgctaagct	cgcactgatt	1080
ttttacctga	gtaggcctag	aaataaacat	gctagctttt	attccagttc	taaccaaaaa	1140
aataaacctt	cgttccacag	aagntgccat	caagtatttc	ctcacgcaag	caaccgcac	1200
cataatcctt	ctaatagcta	tccntttcaa	caatatactc	tccggacaat	gaaccatwac	1260
caataccacc	aatcaatact	catcattaat	aatcataatg	gctatagcaa	taaaactagg	1320
aatagcccc	tttcacttct	gagtcccaga	ggttacccaa	ggcacccttc	tgacatccgg	1380
cctgctcctt	ctcacatgac	aaaaactagc	ccccatctca	atcatatacc	aaatttctcc	1440
ctcattaaac	gtaagccttc	tcctcactct	ttcaatctta	tccatcatgg	caggcagttg	1500
aggtggatta	aaccaaacc	aactacgcaa	aatcttagca	tactcctcaa	ttaccacat	1560
aggatgaata	acagcagttc	taccgtacaa	ccctaacata	accattctta	atttaactat	1620
ttatattatc	ctaactacta	ccgcattcct	actactcaac	ttaaactcca	gcaccacaac	1680
cctactacta	tctcgcacct	gaaacaagct	aacatgacta	acacccttaa	ttccatccac	1740
cctcctctcc	ctaggaggcc	tgcccccgct	aaccggcttt	ttgcccacaa	gggccattat	1800
cgaagaattc	acaaaaaaca	atagcctcat	catccccacc	atcatagcca	tcataccctt	1860
ccttaacctc	tacttctacc	tgcgccta	ctactccacc	tcaatcacac	tactccctat	1920
atctaacaac	gtaaaaataa	aatgacagtt	tgaacacaca	aaacccacc	cattcctccc	1980
cacactcatc	gcccttacca	cactgctcct	acctatctcc	ccttttatgc	taataatctt	2040
atagaaattt	aggttaaata	cagaccaaga	gccttcaaag	ccctcagtaa	gttgcaatac	2100
ttaatttctg	naacagctaa	ggactgcaaa	acccactctt	gcatcaactg	nanagcaaatc	2160
agccacttta	attaagctaa	gcccttacta	gaccaatggg	acttaaacc	acaaacactt	2220
agttaacagc	taagcaccct	aatcaactgg	cttcaatcta	cttctccgc	cgccgggaaa	2280
aaaggcgagg	gaagccccgg	caggtttgaa	gctgcttctt	cgaatttgca	attcaatatg	2340
aaaatcacct	canagctggt	aaaaagaggc	actacccttg	tcttttagatt	tacagtccaa	2400
tgcttcactc	agccatttfa	cctcaccccc	ctttaggttc	gccgaccgtt	gactattctc	2460
tacaaaccac	aaagacattg	gaacactata	cctattatct	ggcgcatgag	ctggagtcct	2520
aggcacagct	ctaagcctcc	ttattcgagc	cgaactgggc	cagccaggca	accttctagg	2580
taacgaccac	atctacaacg	ttatcgtcac	agcccatgca	tttgtaataa	tcttctcat	2640
agtaataccc	atcataatcg	gaggytttg	caastgacta	gttcccctaa	taatcgggtc	2700
ccccgatatg	gsgtttcccc	gcataaaca	cataagcttn	tgactcttac	ccccctctct	2760
cntactcctg	ttngcatctg	ctatagtga	ggccggcgca	ggaacagggt	gaacagtnta	2820
ccctcccttg	gcagggaact	actcccaccc	tggagcctcc	gtagacctaa	ccatcttctc	2880
cctacaccta	gcagggtatct	cctctatctt	agggaccatc	aatttcatca	caacaattat	2940
taataataaa	ccccctgcc	taacccaata	ccaaacgccc	cttttcgtct	gatccgtcct	3000
aatcacagca	gtcttacttc	tcctatctct	cccagtccta	gccgctggca	tactataact	3060
actaacagac	cgtaacctca	acaccacctt	cttcgaccca	gccggaggag	gagaccccat	3120
tctataccaa	cacctattct	gatttttcgg	tcaccctgaa	gtttatattc	tcatacctacc	3180
aggcttcgga	ataatctccc	atattgtaac	ttactactcc	ggaaaaaaag	aaccatttgg	3240
atacataggt	atggtctgag	ctatgatatc	aataggcttc	ctagggttta	tcgtgtgagc	3300
acaccatata	tttacagtag	gaatagacgt	agacacacga	gcatatttca	cctccgctac	3360
cataatcatc	gctatcccca	cggcgctcaa	agtatttagc	tgactcgcca	cactccacgg	3420
aagcaatatg	aatgatctg	ctgcagtgct	ctgagcccta	ggatttattt	ttcttttcac	3480

8/42

cgtagggtggc	ctgactggca	ttgtattagc	aaactcatca	ctagacatcg	tactacacga	3540
cacgtactac	gttgtagccc	acttcacta	tgtcctatca	ataggagctg	tatttgccat	3600
cataggaggc	ttcattcact	gatttcccct	attctcaggc	tacaccctag	accaaaccta	3660
cgccaaaatc	catttcgcta	tcatattcat	cggcgtaaat	ctaactttct	tcccacaaca	3720
citttctcggc	ctatccggaa	tgccccgacg	ttactcggac	tatccccgatg	catacaccac	3780
atgaaatatc	ctatcatctg	taggctcatt	cattttctcta	acagcagtaa	tattaataat	3840
tttcataaatt	tgagaagcct	tcgcttcgaa	gcgaaaagtc	ctaatagtag	aagaaccctc	3900
cataaacctg	gagtgactat	atggatgccc	cccaccctac	cacacattcg	aagaaccctg	3960
atacataaaa	tctagacaaa	aaaggaagga	atcgaacccc	ccaaagctgg	tttcaagcca	4020
accccatggc	ctccatgact	ttttcaaaaa	gatattagaa	aaaccatttc	ataactttgt	4080
caaaagttaa	ttataggcta	aatcctatat	atcttaatgg	cacatgcagc	gcaagtaggt	4140
ctacaagacg	ctacttcccc	tatcatagaa	gagcttatca	tctttcatga	tcacgcccctc	4200
ataatcattt	tccttatctg	cttcctagtc	ctgtacgccc	ttttcctaac	actcacaaca	4260
aaactaacta	atactaactat	ctcagacgct	caggaaatag	aaaccgtctg	aactatcctg	4320
cccgccatca	tcctagtctt	tatcgccctc	ccatncctac	gcctccttta	cataacagac	4380
gagggtcaacg	atccctcctt	taccatcaaa	tcaattggcc	atcaatggta	ctgaacctac	4440
gaataacaccg	actacggcgg	actaatcttc	aactcctaca	tacttcccc	attattccta	4500
gaaccaggcg	acctgcgact	ccttgacggt	gacaatcgag	tagtactccc	ggttgaagcc	4560
cccattcgta	taataattac	atcacaagac	gtcttacact	catgagctgt	ccccacatta	4620
ggcttaaaaa	cagatgcaat	tcccgagcgt	ctaaacccaa	ccactttcac	tgctacacga	4680
ccaggggtat	actacggcca	atgctctgaa	atctgtggag	caaaccagtt	ttatgcccat	4740
cgctctagaa	ttaattcccc	taaaaaatctt	tgaaataggg	cccgtattta	ccctatagca	4800
ccccctctac	cccctctaga	gcccactgta	aagctaacct	agcattaacc	ttttaagtta	4860
aagattaaga	gaaccaacac	ctctttacag	tgaaatgccc	caactaaata	ctaccgtatg	4920
acccaccata	attaccccca	tactccttac	actattcctc	atcacccaac	taaaaaatatt	4980
aaatacaaat	taccacctac	ctccctcacc	aaagcccata	aaaataaaaa	actataacaa	5040
accctgagaa	ccaaaatgaa	cgaaaatctg	ttcacttcat	tcattgcccc	cacaatccta	5100
ggcctacccg	ccgcagtaact	gatcattcta	tttyccccctc	tattgatccc	cacctccaaa	5160
tatctcatca	acaaccgact	aattaccacc	caacaatgac	taatccaact	aacctcaaaa	5220
caaatgatag	ccatacacaa	cactaaggga	cgaaactgat	ctcttatact	agtatcctta	5280
atcattttta	ttgccacaac	taacctcctc	ggactcctgc	ctcactcatt	tacaccaacc	5340
acccaactat	ctataaacct	agccatggcc	atcccccttat	gagcgggcgc	agtgattata	5400
ggctttcgct	ctaagattaa	aaatgcccta	gcccacttct	taccacaagg	cacacctaca	5460
ccccttatcc	ctatactagt	tattatcgaa	accatcgacc	tactcattca	accaatagcc	5520
ctggccgtan	gnctaaccgc	taacattact	gcaggccacc	tactcatgna	cctaattgga	5580
agcgccacac	tagcaataatc	aantattaac	cttccctcta	cacttatcat	tttcacaatt	5640
ctaattctac	tgactatcct	agaaatcgct	gtcgccttaa	tccaagccta	cgtttttaca	5700
cttttagtaa	gcctctacct	gcacgacaac	acataatgac	ccaccaatca	catgcctatc	5760
atatagtaaa	acccagccca	tggcccctaa	caggggccct	ctcanccctc	ctaatagacct	5820
ccggcctagc	catgtgattt	cacttccact	ccacaaccct	ccttatacta	ggcctactaa	5880
ccaacacact	aaccatatac	caatgatggc	gcgatgtaac	acgagaaaagc	acataccaag	5940
gccaccacac	accacctgtc	cagaaaaggc	ttcgatacgg	gataatccta	tttattacct	6000
cccccaact	agggggacac	tggcccccaa	caggcatcac	cccgtaaaat	cccctagaag	6120
tcccactcct	aaacacatcc	gtattactcg	catcagggggt	atcaatcacc	tgagctcacc	6180
atagtctaatt	agaaaacaac	cgaaaacaaa	taattcaagc	actgcttatt	acaattttac	6240
tgggtctcta	ttttaccctc	ctacaagcct	cagngtactt	cgagggtaaa	atattagata	6300
tttcccctga	tacagggtc	aatctttttc	tttttaaagc	aatatttctc	aaagtacttt	6360
tcacagaact	taagtttcat	taagcacttc	actaaaagna	aaagtctgtg	atctaataaa	6420
tttgaaaaat	attgagaatt	agagccccct	cttagatatg	tactgtagct	actcagcttg	6480
ttacagatgg	aagtaaacat	tgtaatattc	acccagcttt	tgagtggatg	tctattaaca	6540
tcacccaaat	gagtattcca	tggaatgcac	tttgcaaaaa	cctattattc	aagaaaaaatt	6600
ctggagcatg	gaaagctatt	aatggataaa	cccattcaca	aaatcacacc	aaatatctaa	6660
aatcatgttt	aaaaatctcct	agaaatgggt	t			6691

Fig. 3B

*Fig. 4A*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
COI	5.8 kb pg	6023	G	A
		6221	T	C
		6242	C	T
		6266	A	C
		6299	A	G
		6366	G	A
		6383	G	A
		6410	C	T
		6452	C	T
		6483	C	T
		6512	T	C
		6542	C	T
		6569	C	A
		6641	T	C
		6935	C	T
		6938	C	T
		7146	A	G
		7232	C	T
		7256	C	T
		7316	G	A
COI	11	6023	G	A
COI	12	6221	T	C
		6242	C	T
		6266	A	C
		6299	A	G
		6366	G	A
		6383	G	A
		6410	C	T
COI	12	6160	T	A
		6182	G	A
		6185	T	C
		5216	T	C
		6221	T	C
		6224	C	T
		6236	C	T
		6242	C	T
		6251	T	C
		6260	G	A
		6266	A	C
		6269	A	C
		6281	A	G
		6326	C	T
		6335	C	T
		6353	A	G
		6356	C	T
		6365	T	C
		6366	G	A
		6378	T	C

*Fig. 4B*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
COI	12	6383	G	A
		6389	C	T
		6392	T	C
		6398	C	T
		6407	T	C
		6410	C	T
		6260	G	A
COI	12	6266	A	C
		6269	A	C
		6281	A	G
		6326	C	T
		6335	C	T
		6353	A	T
		6356	C	T
		6365	T	C
		6366	G	A
		6378	T	C
		6383	G	A
		6392	T	C
		6398	C	T
COI	12	6182	G	A
		6216	T	C
		6221	T	C
		6260	G	A
		6266	A	C
		6269	A	C
		6281	A	G
		6326	C	T
		6335	C	T
		6353	A	T
		6356	C	T
		6365	T	C
		6366	G	A
		6378	T	C
		6383	G	A
		6392	T	C
		6398	C	T
COI	12	6160	T	A
		6182	G	A
		6185	T	C
		5216	T	C
		6221	T	C
		6224	C	T
		6236	C	T
		6242	C	T
		6251	T	C
		6266	A	C
		6269	A	C

*Fig. 4C*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
COI	12	6326	C	T
		6335	C	T
		6353	A	G
		6356	C	T
		6365	T	C
		6366	G	A
		6378	T	C
		6383	G	A
		6392	T	C
		6160	T	A
		6182	G	A
		6185	T	A
COI	12	5216	T	C
		6221	T	C
		6224	C	T
		6242	C	T
		6251	T	C
		6260	G	A
		6266	A	C
		6269	A	C
		6281	A	C
		6326	C	G
		6335	C	T
		6353	A	G
		6356	C	T
		6365	T	C
		6366	G	A
		6378	T	C
		6383	G	A
		6389	C	T
		6392	C	T
		6398	C	T
		6407	T	C
		6410	C	T
		6251	T	C
		6260	G	A
		6266	A	C
		6269	A	C
		6281	A	C
		6326	C	G
		6335	C	T
		6353	A	G
		6356	C	T
		6365	T	C
		6366	G	A
		6378	T	C
		6383	G	A
		6392	T	C

*Fig. 4D*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
COI	12	6398	C	T
		6185	T	C
		6251	T	C
		6260	G	A
		6266	A	C
		6269	A	C
		6281	A	G
		6326	C	T
		6335	C	T
		6353	A	G
		6356	C	T
		6365	T	C
		6366	G	A
		6378	T	A
		6383	G	A
		6392	T	C
COI	12	6398	C	T
		6185	T	C
		6236	T	C
		6251	T	C
		6260	G	A
		6266	A	C
		6269	A	C
		6281	A	G
		6326	C	T
		6335	C	T
		6353	A	G
		6356	C	T
		6365	T	C
		6366	G	A
		6378	T	A
		6383	G	A
COI	12	6392	T	C
		6398	C	T
		6236	C	T
		6251	T	C
		6260	G	A
		6266	A	C
		6269	A	C
		6281	A	G
		6326	C	T
		6335	C	T
		6353	A	G
		6356	C	T
		6365	T	C
		6366	G	A
		6378	T	A
		6383	G	A

*Fig. 4E*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
COI	12	6392	T	C
		6398	C	T
		6236	C	T
		6251	T	C
		6260	G	A
		6266	A	C
		6269	A	C
		6281	A	G
		6326	C	T
		6335	C	T
		6353	A	G
		6356	C	T
		6365	T	C
		6366	G	A
COI	12	6378	T	C
		6383	G	A
		6392	T	C
		6185	T	C
		6236	C	T
		6251	T	C
		6260	G	A
		6266	A	C
		6269	A	C
		6281	A	G
		6326	C	T
		6353	A	G
		6356	C	T
		6365	T	C
COI	12	6366	G	A
		6383	G	A
		6392	T	C
		6398	C	T
		6185	T	C
		6236	C	T
		6251	T	C
		6260	G	A
		6266	A	C
		6269	A	C
		6281	A	G
		6326	C	T
		6353	A	G
		6356	C	T
COI	12	6365	T	C
		6366	G	A
		6383	G	A
		6185	T	C
COI	12	6251	T	C
		6260	G	A
		6266	G	A

*Fig. 4F*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
COI	12	6266	A	C
		6269	A	C
		6281	A	G
		6326	C	T
		6335	C	T
		6353	A	G
		6356	C	T
		6365	T	C
		6366	G	A
		6383	G	A
		6392	T	C
		6398	C	T
COI	12	6185	T	C
		6251	T	C
		6260	G	A
		6266	A	C
		6269	A	C
		6281	A	G
		6326	C	T
		6335	C	T
		6353	A	G
		6356	C	T
		6365	T	C
		6366	G	A
COI	13	6392	T	C
		6398	C	T
		6452	C	T
		6483	C	T
		6512	C	T
		6542	C	T
		6569	C	T
		6641	T	A
		6935	C	T
		6938	C	T
COI	15	7146	A	G
		6950	C	A
COI	15	6962	G	A
		6965	T	G
		6975	T	C
		6990	C	T
		7013	G	A
		7040	T	C
		7064	T	C
		7076	A	G
		7145	A	G
		7146	A	G
		7160	C	A
		7175	T	C



*Fig. 4G*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
COI	15	6950	C	A
		6962	G	A
		6965	T	G
		6975	T	C
		7013	G	A
		7040	T	C
		7064	T	C
		7076	A	G
		7145	C	T
		7146	A	G
		7160	C	A
		7175	T	C
COI	15	6932	A	G
		6935	C	T
		6938	C	T
		6944	T	C
		6950	C	A
		6962	G	A
		6965	T	G
		6975	T	C
		6990	C	T
		7013	G	A
		7022	T	C
		7028	C	T
		7037	C	T
		7040	T	C
		7064	T	C
		7072	T	C
		7079	C	T
		7100	A	G
		7112	C	T
		7145	C	T
COI	15	7146	A	G
		7160	C	A
		7169	T	C
		7175	T	C
		6932	A	G
		6935	C	T
		6938	C	T
		6944	T	C
		6950	C	A
		6962	G	A
		6965	T	G
		6975	T	C
		6990	C	T
		7013	G	A
		7022	T	C
		7028	C	T

*Fig. 4H*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
COI	15	7037	C	T
		7064	T	C
		7072	T	C
		7079	C	T
		7100	A	G
		7112	C	T
		7145	C	T
		7146	A	G
		7160	C	A
		7169	T	C
		7175	T	C
		6929	A	G
		6938	C	T
COI	15	6944	T	C
		6950	C	A
		6956	T	C
		6962	G	A
		7013	G	A
		7022	T	C
		7040	T	C
		7064	T	C
		7076	A	G
		7133	C	T
		7145	C	T
		7146	A	G
		7169	T	C
COI	15	6929	A	G
		6938	C	T
		6944	T	C
		6950	C	A
		6956	T	C
		7013	G	A
		7022	T	C
		7040	T	C
		7064	T	C
		7076	A	G
		7133	C	T
		7145	C	T
		7146	A	G
COI	15	7169	T	C
		6929	A	G
		6938	C	T
		6944	T	C
		6950	C	A
		6956	T	C
		6962	G	A
		7013	G	A
		7022	T	C

*Fig. 4I*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
		7040	T	C
		7064	T	C
		7076	A	G
		7133	C	T
		7146	A	G
		7169	T	C
COI	15	6950	C	A
		6962	G	A
		6965	T	G
		6975	T	C
		6990	C	T
		7013	G	A
		7040	T	C
		7064	T	C
		7076	A	G
		7145	C	T
		7146	A	G
		7160	C	A
		7175	T	C
COI	15	6932	A	G
		6938	C	T
		6950	C	A
		6962	G	A
		6965	T	G
		6975	T	C
		6990	C	T
		7013	G	A
		7022	T	C
		7028	C	T
		7037	C	T
		7040	T	C
		7064	T	C
		7072	T	C
		7079	C	T
		7100	A	G
		7112	C	T
		7146	A	G
		7160	C	A
		7169	T	C
		7175	T	C
COI	16	7232	C	T
		7256	C	T
		7316	G	A
COII	5.8 kb pg	7650	C	T
		7705	T	C
		7810	C	T
		7868	C	T
		7891	C	T

*Fig. 4J*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution, (or Deletion)
		7912	G	A
		8021	A	G
		8065	G	A
		8140	C	T
		8152	G	A
		8167	T	C
		8196-8197	AC	deletion
		8203	C	T
COII	21/22/23	7650	C	T
		7705	T	C
		7810	C	T
		7868	C	T
		7891	C	T
		7912	G	A
		8021	A	G
		8065	G	A
		8140	C	T
		8152	G	A
		8167	T	C
		8196-8197	AC	deletion
		8203	C	T
COII	21/22/23	7650	C	T
		7705	T	C
		7810	C	T
		7868	C	T
		7891	C	T
		7912	G	A
		8021	A	G
COII	21	7650	C	T
		7705	T	C
		7810	C	T
COII	21	7650	C	T
COII	21/22/23	7650	C	T
		7705	T	C
		7861	T	C
		7868	C	T
		7891	C	T
		7900	C	T
		7912	G	A
		7927	C	T
		8011	A	G
		8021	A	G
		8038	T	C
COII	21/22/23	7650	C	T
		7663	C	T
		7705	T	C
		7810	C	T
		7868	C	T

*Fig. 4K*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
		7891	C	T
		7912	G	A
		8021	A	G
COII	21/22/23	7650	C	T
		7705	T	C
		7810	C	T
		7861	T	C
		7868	C	T
		7891	C	T
		7900	C	T
		7912	G	A
		7927	C	T
		8011	A	G
		8021	A	G
COII	22	7868	C	T
		7891	C	T
		7912	G	A
COII	21/22/23	7650	C	T
		7705	T	C
		7757	G	A
		7810	C	T
		7868	C	T
		7891	C	T
		7912	G	A
		8021	A	G
COII	22	7868	C	T
		7891	C	T
		8021	A	G
COII	21/22	7650	C	T
		7810	C	T
		7868	C	T
		7891	C	T
		7912	G	A
		8021	A	G
COII	21/22	7650	C	T
		7705	T	C
		7810	C	T
		7861	T	C
		7868	C	T
		7891	C	T
		7900	C	T
		7912	G	A
		7927	C	T
		8011	A	G
		8021	A	G
		8038	T	C
COII	21	7650	C	T
		7810	C	T

*Fig. 4L*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
COIII	5.8 kb pg	9254	A	G
		9325	T	C
		9329	G	C
		9335	C	T
		9434	A	G
		9540	T	C
		9545	A	G
		9548	G	A
		9559	G	C
		9629	A	G
ATPase 8	26/5.8 kb pg	8392	G	A
		8455	C	T
		8461	C	T
		8503	T	C
		8545	G	A
ATPase 8/6	519 bp pg	8541	G	A
		8557	G	A
		8562	C	T
		8566	A	G
		8568	C	A
ATPase 8	26	8371	C	A
		8374	A	G
		8380	T	C
		8383	T	C
		8386	C	T
		8392	G	A
		8401	A	C
		8410	C	A
		8419	T	C
		8428	C	T
		8455	C	T
		8461	C	T
		8467	C	T
		8468	C	T
		8469	T	C
		8473	T	deletion
		8476	C	T
		8485	G	A
		8491	A	T
		8503	T	C
		8506	T	C
		8507	A	G
		8509	C	T
		8541	G	A
		8558	C	G
		8562	C	T
		8566	A	G

*Fig. 4M*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
ATPase 8	26/519 bp pg	8568	C	A
		8371	C	A
		8374	A	G
		8383	T	C
		8386	C	T
		8392	G	A
		8395	C	T
		8396	A	G
		8398	C	T
		8401	A	C
		8404	T	C
		8410	C	A
		8419	T	C
		8422	A	G
		8423	C	T
		8428	C	T
		8450	T	C
		8459	A	C
		8463	A	G
		8467	A	T
		8470	C	G
		8473	A	C
		8474	C	A
		8485	G	A
		8486	C	T
		8487	C	T
		8488	C	T
		8491	A	T
		8503	T	C
		8506	T	C
		8508	A	G
		8509	C	T
		8512	A	G
		8539	C	T
		8541	G	A
		8557	G	A
		8562	C	T
		8566	A	G
		8568	C	A
ATPase 6	5.8 kb pg	8545	G	A
		8655	C	T
		8677	A	C
		8701	A	G
		8718	A	G
		8860	A	G
		8943	C	T
		9060	C	A
		9075	C	T

*Fig. 4N*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution, (or Deletion)
ATPase 6	519 bp pg	9103	C	T
		9168	C	T
		9175	C	T
		8584	G	A
		8591	T	C
		8592	G	A
		8598	T	C
		8610	T	C
		8611	C	T
		8614	T	C
		8617	A	G
		8622	C	A
		8634	T	C
		8661	C	T
		8674	A	C
		8676	C	T
		8677	A	C
		8682	A	C
		8687	G	A
		8697	C	T
		8703	C	G
		8709	C	T
		8714	C	T
		8718	A	G
		8730	A	G
		8733	T	C
		8743	G	A
		8745	A	T
		8749	T	C
		8751	A	G
		8754	C	T
		8775	C	T
		8788	C	T
		8793	T	C
		8810	C	T
ND1	63/5.8 kb pg	4048	G	A
		4104	A	G
ND2	5.8 kb pg	4496	C	T
		4736	T	C
		4769	A	G
		4856	T	C
		4904	C	T
		4914	C	T
		4940	C	T
		4958	A	G
		4985	G	A
		4991	G	A
		5041	T	C



*Fig. 40*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
ND2	72	5147	G	A
		5320	C	T
		5351	A	G
		5387	C	T
		5426	T	C
		5471	G	A
		5474	A	G
		5498	A	G
		4736	T	C
		4856	T	C
ND2	73	4904	C	T
		4914	C	T
		4940	C	T
		4958	A	G
		4991	G	A
		5041	T	C
		5041	T	C
ND2	74	5147	G	A
		5320	C	T
		5351	A	G
		5387	C	T
		5426	T	C
		5471	G	A
		5474	A	G
ND3	75	5498	A	G
		10040	C	T
		10041	A	G
		10043	C	T
		10071	T	C
		10077	T	C
		10083	A	G
		10088	C	T
		10101	T	C
		10107	C	T
		10113	A	G
		10124	T	C
		10128	C	T
		10143	G	A
		10145	C	T
		10162	C	A
		10172	G	A
		10175	C	T
		10187	T	C
		10188	A	C
		10192	C	T
		10197	G	A
		10203	G	A
		10205	C	T

*Fig. 4P*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
		10217	A	G
		10235	T	C
		10238	T	C
		10244	C	T
		10245	T	C
		10256	T	C
		10268	C	T
		10274	T	A
		10275	T	A
		10277	A	C
		10281	C	T
		10304	T	C
		10308	C	T
		10309	T	A
		10310	G	A
		10318	T	A
		10322	T	C
		10325	G	A
		10346	C	T
		10353	G	A
		10361	T	C
		10364	G	A
		10373	G	A
		10377	C	T
		10378	T	C
		10385	A	G
		10398	A	G
		10410	T	C
		10421	C	T
ND3	75	10007	T	G
		10008	A	C
		10009	G	T
		10037	T	C
		10043	C	A
		10044	A	G
		10046	T	C
		10047	C	T
		10048	A	G
		10061	A	T
		10065	A	T
		10067	T	C
		10070	C	A
		10071	C	A
		10074	T	G
		10075	A	C
		10076	T	C
		10077	T	C
		10084	T	C

*Fig. 4Q*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
		10085	C	T
		10086	A	G
		10091	C	T
		10092	C	T
		10094	C	A
		10095	C	T
		10098	G	A
		10100	C	T
		10101	T	C
		10104	C	T
		10107	C	T
		10110	A	G
		10113	A	G
		10115	T	A
		10130	A	T
		10131	C	G
		10134	C	T
		10139	C	T
		10142	C	T
		10143	G	A
		10144	G	C
		10145	C	T
		10162	C	G
		10165	C	T
		10169	C	T
		10172	G	A
		10175	C	T
		10176	G	A
		10178	C	A
		10181	C	T
		10184	C	T
		10187	T	A
		10188	A	T
		10191	T	A
		10194	C	T
		10196	C	T
		10197	G	A
		10201	G	A
		10203	G	C
		10207	C	T
		10208	T	C
		10226	C	T
		10227	T	C
		10235	T	C
		10238	T	C
		10241	C	A
		10244	C	T
		10245	T	C

*Fig. 4R*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
		10247	A	C
		10248	T	C
		10253	T	C
		10257	C	deletion
		10265	T	C
		10267	C	T
		10271	C	A
		10272	C	T
		10274	T	C
		10275	T	C
		10283	A	G
		10295	A	T
		10303	C	A
		10310	G	T
		10311	C	A
		10312	C	T
		10320	G	A
		10322	T	C
		10324	T	G
		10325	G	T
		10326	T	A
		10329	T	A
		10334	C	T
		10335	T	C
		10338	T	C
		10343	C	T
		10345	T	C
		10349	C	T
		10353	G	A
		10354	C	T
		10359	A	G
		10361	T	G
		10364	G	A
		10365	G	A
		10366	C	T
		10367	C	T
		10373	G	A
		10377	C	A
		10378	T	C
		10379	A	C
		10394	C	T
		10398	A	G
		10399	C	T
		10400	C	T
		10404	T	C
		10410	T	A
		10421	C	T
		10422	A	T

*Fig. 4S*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
		10426	C	deletion
		10427	G	deletion
		10428	A	deletion
		10429	A	deletion
ND3	75	10040	C	T
		10041	A	G
		10043	C	T
		10071	T	C
		10077	T	C
		10083	A	G
		10088	C	T
		10101	T	C
		10107	C	T
		10113	A	G
		10124	T	C
		10128	C	T
		10143	G	A
		10145	C	T
		10162	C	A
		10172	G	A
		10175	C	T
		10187	T	C
		10188	A	C
		10192	C	T
		10197	G	A
		10203	G	A
		10205	C	T
		10217	A	G
		10235	T	C
		10238	T	C
		10244	T	C
		10245	T	C
		10256	T	C
		10268	C	T
		10274	T	A
		10275	T	C
		10277	A	G
		10281	C	T
		10304	T	C
		10308	C	T
		10310	G	A
		10318	T	C
		10322	T	C
		10325	G	A
		10336	T	A
		10346	C	T
		10353	G	A
		10361	T	C

*Fig. 4T*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
		10364	G	A
		10370	T	C
		10373	G	A
		10377	C	T
		10378	T	C
		10385	A	G
		10398	A	G
		10410	T	C
		10421	C	T
ND3	75	10040	C	T
		10041	A	G
		10043	C	T
		10071	T	C
		10083	A	G
		10088	C	T
		10101	T	C
		10107	C	T
		10113	A	G
		10124	T	C
		10128	C	T
		10143	G	A
		10145	C	T
		10162	C	A
		10172	G	A
		10175	C	T
		10187	T	C
		10188	A	C
		10192	C	T
		10197	G	A
		10203	G	A
		10205	C	T
		10217	A	G
		10235	T	C
		10238	T	C
		10244	C	T
		10245	T	C
		10256	T	C
		10268	C	T
		10274	T	A
		10275	T	C
		10277	A	G
		10281	C	T
		10304	T	C
		10308	C	T
		10310	G	A
		10318	T	C
		10322	T	C
		10325	G	A

*Fig. 4U*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
ND4L	81	10346	C	T
		10353	G	A
		10361	T	C
		10364	G	A
		10370	T	C
		10373	G	A
		10377	C	T
		10378	T	C
		10385	A	G
		10398	A	G
		10410	T	C
		10421	C	T
		10475	C	T
		10501	C	A
		10547	C	G
		10586	G	A
		10589	G	A
		10601	T	C
		10646	G	A
ND4	82	10652	T	C
		10670	C	T
		10677	G	A
		10679	G	A
		10685	A	G
		10688	G	A
		10721	A	G
		10750	A	G
		10750	A	G
		10774	C	T
		10785	T	C
		10810	T	C
		10846	C	T
		10866	T	C
		10873	T	C
ND4	82	10885	T	C
		10915	T	C
		10919	T	C
		10920	C	T
		10922	A	T
		10927	T	C
		10945	A	G
		11016	G	A
		10750	A	G
		10774	C	T
		10810	T	C
		10846	C	T
		10866	T	C
		10873	T	C

*Fig. 4V*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
ND4	82	10885	T	C
		10915	T	C
		10919	C	T
		10920	T	C
		10922	A	T
		10927	T	C
		10945	A	G
		11016	G	A
		10750	A	G
		10774	C	T
		10786	T	C
		10800	T	A
		10808	C	T
		10810	T	C
		10822	C	T
		10846	C	T
ND4	82	10866	T	C
		10873	T	C
		10915	T	C
		10920	T	C
		10922	A	T
		10927	T	C
		10945	A	G
		10975	C	T
		11002	A	G
		11009	T	C
		11016	G	A
		11017	T	C
		11053	A	G
		10750	A	G
		10775	G	A
		10786	T	C
		10801	G	A
		10808	C	T
		10810	T	C
		10822	C	T
		10846	C	T
		10866	T	C
		10873	T	C
		10915	T	C
		10920	T	C
		10922	A	T
		10927	T	C
		10945	A	G
		10975	C	T
		11002	A	G
		11009	T	C
		11016	G	A



*Fig. 4W*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
ND4	83	11017	T	C
		11053	A	G
		11083	A	G
		11097	C	T
		11147	T	C
		11176	G	A
		11197	C	T
		11233	T	C
		11254	T	C
		11260	T	C
		11281	A	G
		11284	C	T
		11291	C	T
		11302	C	T
ND4	85	11335	T	C
		11590	A	G
		11662	T	C
		11708	A	G
		11719	G	A
		11767	C	T
		11770	T	C
		11788	C	T
		11809	T	C
		11827	T	C
		11852	G	A
		11857	C	T
		11887	G	A
ND4	86	11857	C	T
		11887	G	A
		11914	G	A
		11963	G	A
		12007	G	A
		12013	A	G
		12018	C	G
		12064	C	T
		12088	C	T
		12091	T	C
		12115	C	T
ND4	86	12136	T	C
		11854	T	C
		11863	C	T
		11886	A	G
		11875	T	C
		11887	G	A
		11898	C	T
		11899	T	C
		11902	G	A
		11908	A	G

*Fig. 4X*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution, (or Deletion)
		11912	A	T
		11914	G	A
		11929	T	C
		11935	T	C
		11944	T	C
		11953	C	T
		11963	G	A
		11986	C	T
		12007	G	A
		12013	A	G
		12019	C	A
		12047	T	C
		12049	C	T
		12064	C	T
		12071	G	A
		12091	T	C
		12106	C	T
		12112	C	T
		12115	C	T
		12131	T	A
		12136	T	C
ND5	93	12346	C	T
		12348	C	T
		12349	A	G
		12351	T	C
		12354	T	C
		12362	C	T
		12367	A	G
		12372	G	A
		12379	C	T
		12394	C	A
		12397	A	G
		12403	C	T
		12406	G	A
		12408	T	C
		12423	A	G
		12425	A	G
		12426	C	T
		12438	T	C
		12450	C	A
		12454	G	A
		12456	C	A
		12463	A	T
		12474	C	T
		12477	T	C
		12481	T	C
		12501	G	A
		12528	G	A

*Fig. 4Y*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
ND5	93	12543	C	A
		12557	C	T
		12558	C	A
		12561	G	A
		12567	C	G
		12574	T	G
		12576	C	T
		12603	C	A
		12616	T	C
		12618	G	A
		12624	T	C
		12630	G	A
		12651	G	A
		12654	A	G
		12662	A	G
		12346	C	T
		12349	A	G
		12358	A	C
		12367	A	G
		12372	G	A
		12379	C	T
		12390	C	T
		12406	G	A
		12417	C	T
		12432	C	T
		12441	T	A
		12454	G	A
		12466	T	C
		12469	A	G
		12474	C	T
		12501	G	A
		12503	G	A
		12519	T	C
		12528	G	A
		12540	A	G
		12543	C	A
		12561	G	A
		12585	C	T
		12588	C	T
		12603	C	T
		12616	T	C
		12627	A	G
		12630	G	A
		12651	G	A
		12662	A	G
ND5	93	12302	C	A
		12303	C	A
		12311	T	C

*Fig. 4Z*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution, (or Deletion)
		12312	T	A
		12316	G	A
		12319	C	T
		12338	T	C
		12341	C	A
		12346	C	T
		12348	C	T
		12349	A	deletion
		12352	A	T
		12354	T	C
		12357	A	G
		12358	A	G
		12360	C	T
		12362	C	T
		12363	C	A
		12364	C	A
		12365	T	C
		12367	A	G
		12372	G	A
		12374	C	T
		12375	T	C
		12376	T	C
		12379	C	T
		12387	C	T
		12388	C	A
		12391	A	G
		12392	T	A
		12393	C	T
		12394	C	T
		12395	T	A
		12396	T	C
		12397	A	T
		12398	C	A
		12399	C	T
		12400	A	T
		12401	C	T
		12405	C	A
		12406	G	T
		12409	A	C
		12411	C	A
		12413	C	T
		12414	T	C
		12415	A	C
		12416	A	T
		21417	C	G
		12418	A	C
		12423	A	T
		12425	A	G

*Fig. 4AA*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
		12426	C	G
		12427	T	C
		12428	C	T
		12429	A	C
		12430	T	A
		12431	A	T
		12432	C	A
		12436	C	A
		12438	T	A
		12440	A	T
		12441	T	A
		12442	G	T
		12443	T	G
		12444	A	T
		12448	T	A
		12449	C	A
		12453	T	C
		12454	G	T
		12455	T	A
		12456	C	T
		12457	G	C
		12458	C	G
		12459	A	C
		12460	T	C
		12461	C	A
		12462	C	A
		12463	A	T
		12464	C	G
		12466	T	C
		12469	A	C
		12470	T	A
		12472	A	C
		12473	T	G
		12474	C	T
		12475	A	T
		12476	G	A
		12477	T	G
		12479	T	C
		12480	C	T
		12481	T	C
		12482	T	A
		12483	C	T
		12486	C	deletion
		12487	A	deletion
		12494	T	C
		12495	A	G
		12498	C	T
		12501	G	A

*Fig. 4BB*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
		12505	C	A
		12506	T	C
		12518	T	C
		12519	T	C
		12528	G	A
		12537	C	T
		12541	G	A
		12542	C	T
		12543	C	G
		12545	C	T
		12546	A	G
		12548	C	T
		12552	A	G
		12555	A	T
		12556	A	C
		12557	C	T
		12559	C	A
		12561	G	A
		12566	C	T
		12567	C	A
		12596	T	C
		12599	T	A
		12603	C	T
		12609	T	A
		12614	C	T
		12615	A	G
		12616	T	C
		12618	G	A
		12622	G	A
		12627	A	C
		12630	G	A
		12633	C	T
		12636	C	T
		12637	A	G
		12649	C	A
		12651	G	A
		12663	C	T
ND5	93	12302	C	A
		12303	C	A
		12312	T	A
		12315	G	T
		12322	C	T
		12346	C	T
		12348	C	T
		12349	A	T
		12350	C	T
		12352	A	T
		12354	T	C

*Fig. 4CC*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
		12357	A	T
		12360	C	T
		12362	C	T
		12363	C	A
		12364	C	A
		12370	C	T
		12372	G	A
		12374	C	T
		12375	T	C
		12379	C	T
		12383	T	C
		12384	T	C
		12385	C	T
		12386	C	T
		12387	C	A
		12390	C	A
		12393	C	A
		12394	C	A
		12398	C	T
		12399	C	T
		12400	A	G
		12403	C	T
		12406	G	A
		12407	T	C
		12408	T	C
		12415	A	G
		12416	A	G
		12417	C	T
		12420	A	G
		12423	A	G
		12429	A	G
		12432	C	T
		12435	C	A
		12436	C	T
		12448	T	A
		12449	C	G
		12451	A	deletion
		12452	T	deletion
		12454	G	A
		12457	G	T
		12460	T	A
		12461	C	G
		12465	C	T
		12468	T	C
		12469	A	G
		12470	T	C
		12471	T	C
		12474	C	T

*Fig. 4DD*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution, (or Deletion)
		12477	T	C
		12481	T	C
		12483	C	T
		12486	C	G
		12487	A	G
		12498	C	T
		12504	C	T
		12505	C	A
		12506	T	C
		12518	T	C
		12519	T	C
		12522	T	C
		12528	G	A
		12535	C	T
		12537	C	T
		12541	G	A
		12542	C	T
		12543	C	A
		12548	C	T
		12555	A	C
		12556	A	C
		12558	C	A
		12559	C	A
		12561	G	A
		12567	C	A
		12568	C	T
		12592	T	deletion
		12596	T	deletion
		12600	A	G
		12603	C	T
		12606	C	T
		12609	T	A
		12616	T	C
		12618	G	A
		12621	C	T
		12622	G	A
		12624	T	C
		12627	A	T
		12630	G	A
		12633	C	T
		12636	C	T
		12649	C	deletion
		12651	G	A
		12654	A	G
		12663	C	G
		12669	C	T
ND6	91	14170	A	T
		14172	T	G



*Fig. 4EE*

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]

Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
ND6	91	14182	T	C
		14274	A	G
		14284	C	T
		14305	G	A
		14311	T	C
		14323	G	A
		14362	C	T
		14364	G	A
		14368	C	T
		14377	C	T
		14384	G	A
		14386	T	C
		14404	C	T
		14152	A	G
		14155	C	A
		14165	A	C
		14197	T	G
		14215	T	C
		14218	T	C
		14255	C	T
		14230	A	G
		14237	A	G
		14239	C	T
		14275	C	T
		14276	C	T
		14280	A	G
		14287	T	C
		14305	G	A
		14323	G	A
		14326	T	C
		14334	C	T
		14350	C	T
		14356	C	T
		14362	C	T
		14368	C	T
		14371	T	C
		14374	T	C
		14377	C	T
		14380	C	T
		14384	G	T
		14389	C	T
		14404	C	T
tRNA Lys	519 bp pg	8310	T	C
		8311	T	C
		8336	T	C
		8345	C	T
		8348	A	T
		8349	C	T


40/42

Human exmtDNA Nucleotide Substitutions and Deletions Relative to  
Corresponding Human mtDNA Sequence [SEQ ID NO:2]




Mito- chondrial Gene Region	Fragment	Human mtDNA Nucleotide Position No.	Human mtDNA SEQ ID NO:2	Human exmtDNA Substitution (or Deletion)
tRNA Ile	5.8 kb pg	8351	C	T
		4312	C	T
		4318	C	T
tRNA Gln	5.8 kb pg	4375	C	G
		4382	C	A
		4398	C	T
		4456	C	T
tRNA Met	5.8 kb pg	5821	G	A
tRNA Cys	5.8 kb pg	5840	C	T
tRNA Tyr	5.8 kb pg	7521	G	A

*Fig. 4FF*

41/42

*Fig. 5*

42/42

*Fig. 6A**Fig. 6B**Fig. 6C*

## SEQUENCE LISTING

<110> Mitokor  
 Herrnstadt, Corrina  
 Ghosh, Soumitra S.  
 Clevenger, William  
 Fahy, Eoin F.  
 Davis, Robert E.

<120> DIAGNOSTIC METHOD BASED ON  
 QUANTIFICATION OF EXTRAMITOCHONDRIAL DNA

<130> 660088.416PC

<140> US

<141> 1999-06-14

<160> 112

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 519

<212> DNA

<213> Homo sapien

<400> 1

cactgtaaag ctaaccacgc attaaccttt taagttaaag actaagagaa tcatattctc	60
tttacagtga aatgccacag ctaaatacca ctgtatgacc tgctatcatc accccaatac	120
tcctcacggtt atttctcatc acccaactaa aaataactaaa cacacactgc catctgccc	180
cctcaccaaa atttattaaa ataaaaaact acagtaagcc ctgagaacca aaatgaacga	240
aaattttattc gcttcattca ttacccttac agtactaggc ctaccgcga cagtaccaat	300
catcctattt ccccccttac tgggtcccaac ctccaaatac ctcatcaaca accgactaat	360
caccactcaa caatgactac ttcaactcac cttaaaacaa ataataacga tacataacat	420
taagggacga acctgggtccc ttataactaat ttccctgatt attttttattg ccacaactaa	480
tctcctcgga ctcttgcccc actcatttac accaatcac	519

<210> 2

<211> 16569

<212> DNA

<213> Homo sapien

<400> 2

gatcacaggt ctatcacctt attaacctt cacgggagct ctccatgcat ttggtatttt	60
cgtctggggg gtatgcacgc gatagcattg cgagacgctg gagccggagc accctatgtc	120
gcagtatctg tctttgatcc ctgcctcatc ctattattta tcgcacctac gttcaatatt	180
acaggcgaac atacttacta aagtgtgtta attaatattt gcttgtagga cataataata	240
acaattgaat gtctgcacag ccactttcca cacagacatc ataacaaaaa atttccacca	300
aacccccctt cccccgttc tggccacagc acttaaacac atctctgcca aacccccaaa	360
acaaagaacc ctaacaccag ctaaccaga ttcaaattt tatcttttgg cggtatgcac	420
ttttaacagt cccccccaa ctaacacatt attttccctt cccactccca tactactaat	480
ctcatcaata caacccccgc ccattcctacc cagcacacac acaccgctgc taaccccata	540
ccccgaacca accaaacccc aaagacaccc cccacagttt atgtagctta cctcctcaaa	600

gcaatacact	gaaaaatgttt	agacggggtc	acatcacccc	ataaacaaat	aggtttggtc	660
ctagcctttc	tatttagctct	tagtaagatt	acacatgcaa	gcacccccgt	tccagtgagt	720
tcaccctcta	aatcaccacg	atcaaaaggg	acaagcatca	agcacgcagc	aatgcagctc	780
aaaacgctta	gcctagccac	acccccacgg	gaaacagcag	tgattaacct	ttagcaataa	840
acgaaagttt	aactaagcta	tactaacccc	agggttggtc	aatttcgtgc	cagccaccgc	900
ggtcacacga	ttaacccaag	tcaatagaag	ccggcgtaaa	gagtgtttta	gatcaccccc	960
tccccaataa	agctaaaact	cacctgagtt	gtaaaaaact	ccagttgaca	caaaatagac	1020
tacgaaagtg	gctttaacat	atctgaacac	acaatagcta	agacccaaac	tgggattaga	1080
taccccacta	tgcttagccc	taaacctcaa	cagttaaatc	aacaaaactg	ctcgccagaa	1140
cactacgagc	cacagcttaa	aactcaaagg	acctggcggt	gcttcatatc	cctctagagg	1200
agcctgttct	gtaatcgata	aaccccgatc	aacctcacca	cctcttgctc	agcctatata	1260
ccgccatctt	cagcaaaccc	tgatgaaggc	tacaaagtaa	gcgcaagtac	ccacgtaaag	1320
acgttaggtc	aaggtgtagc	ccatgaggtg	gcaagaaatg	ggctacattt	tctaccccag	1380
aaaactacga	tagcccttat	gaaacttaag	ggtcgaaagt	ggatttagca	gtaaactaag	1440
agtagagtgc	ttagttgaac	agggccctga	agcgcgtaac	caccgcccgt	cacctcctc	1500
aagtataactt	caaaggacat	ttaactaaaa	cccctacgca	tttatataga	ggagacaagt	1560
cgtaacatgg	taagtgtact	ggaaagtgca	cttggacgaa	ccagagtgtg	gcttaacaca	1620
aagcacccaa	cttacactta	ggagatttca	acttaacttg	accgctctga	gctaaaccta	1680
gccccaaacc	cactccacct	tactaccaga	caaccttagc	caaaccattt	acccaaataa	1740
agtataggcg	atagaaattg	aaacctggcg	caatagatat	agtaccgcaa	gggaaagatg	1800
aaaaattata	accaagcata	atatagcaag	gactaacccc	tataccttct	gcataatgaa	1860
ttaactagaa	ataactttgc	aaggagagcc	aaagctaaga	ccccgaaac	cagacgagct	1920
acctaagaac	agctaaaaga	gcacacccgt	ctatgtagca	aaatagtggg	aagatttata	1980
ggtagaggcg	acaaacctac	cgagcctggg	gtagctgggt	tgtccaagat	agaatcttag	2040
ttcaacttta	aattttgccca	cagaacctct	taaatcccc	tgtaaattta	actgttagtc	2100
caaagaggaa	cagctctttg	gacactagga	aaaaaccttg	tagagagagt	aaaaaattta	2160
acaccatag	taggcctaaa	agcagccacc	aattaagaaa	gcgttcaagc	tcaacaccca	2220
ctacctaaaa	aatcccaaac	atataactga	actcctcaca	cccaatttga	ccaatctatc	2280
accctataga	agaactaatg	ttagtataag	taacatgaaa	acattctcct	ccgcataagc	2340
ctgcgtcaga	ttaaaaact	gaactgacaa	ttaacagccc	aatatctaca	atcaaccaac	2400
aagtcattat	tacctcact	gtcaacccaa	cacaggcatg	ctcataagga	aagggttaaaa	2460
aaagtaaaag	gaactcggca	aatcttacct	cgctgtttta	ccaaaaacat	cacctctagc	2520
atcacaagta	ttagaggcac	cgcttgccca	gtgacacatg	tttaacggcc	gcgggtacct	2580
aaccgtgcaa	aggtagcata	atcacttggt	ccttaaatag	ggacctgtat	gaatggctcc	2640
acgaggggtc	agctgtctct	tacttttaac	cagtgaat	gacctgcccc	tgaagaggcg	2700
ggcataaac	agcaagacga	gaagacccta	tggagcttta	atttattaat	gcaaacagta	2760
cctaacaac	ccacaggtcc	taaactacca	aacctgcatt	aaaaatttgc	gttggggcga	2820
cctcggagca	gaacccaacc	tccgagcagt	acatgctaag	acttcaccag	tcaaagcgaa	2880
ctactatact	caattgatcc	aataaactga	ccaacggaac	aagttaccct	agggataaca	2940
gcgcaatcct	attctagagt	ccatatcaac	aatagggttt	acgacctcga	tgttggatca	3000
ggacatccc	atgggtgcagc	cgctattaaa	ggttcgtttg	ttcaacgatt	aaagtcctac	3060
gtgatctgag	ttcagaccgg	agtaatccag	gtcggtttct	atctaccttc	aaattcctcc	3120
ctgtacgaaa	ggacaagaga	aataaggcct	acttcacaaa	gcgccttccc	ccgtaaatga	3180
tatcatctca	acttagtatt	atccccacac	ccacccaaga	acagggtttg	ttaagatggc	3240
agagcccgtt	aatcgcataa	aacttaaaac	tttacagtca	gaggttcaat	tcctcttctt	3300
aacaacatac	ccatggccaa	cctcctactc	ctcattgtac	ccattctaat	cgcaatggca	3360
ttcctaattg	ttaccgaacg	aaaaattcta	ggctatatat	aactacgcaa	aggccccaac	3420
gttgtaggcc	cctacgggct	actacaaccc	ttcgctgacg	ccataaaact	cttcacccaaa	3480
gagcccttaa	aacccgccac	atctaccatc	acctcttaca	tcaccgcccc	gaccttagct	3540
ctcaccatcg	ctcttctact	atgaaccccc	ctccccatac	ccaaccccc	ggtcaacctc	3600
aacctaggcc	tcctatttat	tctagccacc	tctagcctag	ccgtttactc	aatcctctga	3660
tcagggtgag	catcaaactc	aaactacgcc	ctgatcggcg	cactgcgagc	agtagcccaa	3720
acaatctcat	atgaagtcac	cctagccatc	attctactat	caacattact	aataagtggc	3780
tcctttaacc	tctccacct	tatcacaaca	caagaacacc	tctgattact	cctgccatca	3840
tgacccttgg	ccataatatg	atttatctcc	acactagcag	agaccaaccg	aacccccctc	3900

gaccttgccg	aaggggagtc	cgaactagtc	tcaggcttca	acatcgaata	cgccgcaggc	3960
cccttcgccc	tattcttcat	agccgaatac	acaaacatta	ttataataaa	caccctcacc	4020
actacaatct	tcctaggaac	aacatatgac	gcactctccc	ctgaactcta	cacaacatat	4080
tttgtcacca	agaccctact	tctaaccctc	ctgttcttat	gaattcgaac	agcatacccc	4140
cgattccgct	acgaccaact	catacacctc	ctatgaaaaa	acttcctacc	actcacccta	4200
gcattactta	tatgatatgt	ctccataccc	attacaatct	ccagcattcc	ccctcaaacc	4260
taagaaatat	gtctgataaa	agagttactt	tgatagagta	aataatagga	gcttaaacc	4320
ccttatttct	aggactatga	gaatcgaacc	catccctgag	aatccaaaat	tctccgtgcc	4380
acctatcaca	ccccatccta	aagtaaggte	agctaaataa	gctatcgggc	ccataccccg	4440
aaaatgttgg	ttataaccctt	cccgactacta	ttaatcccct	ggcccaaccc	gtcatctact	4500
ctaccatctt	tgcaggcaca	ctcatcacag	cgctaagctc	gcactgattt	tttacctgag	4560
taggcctaga	aataaacatg	ctagctttta	ttccagttct	aacccaaaaa	ataaaccttc	4620
gttccacaga	agctgccatc	aagtatttcc	tcacgcaagc	aaccgcatcc	ataatccttc	4680
taatagctat	cctcttcaac	aataactctt	ccggacaatg	aaccataacc	aataactacca	4740
atcaatactc	atcattaata	atcataatag	ctatagcaat	aaaactagga	atagccccct	4800
ttcacttctg	agtcccagag	gttaccacaag	gcacccctct	gacatccggc	ctgcttcttc	4860
tcacatgaca	aaaactagcc	cccatctcaa	tcataatacca	aatctctccc	tactaaacg	4920
taagccttct	cctcactctc	tcaatcttat	ccatcatagc	aggcagttga	ggtaggattaa	4980
accagaccca	gctacgcaaa	atcttagcat	actcctcaat	taccacata	ggatgaataa	5040
tagcagttct	accgtacaac	cctaacataa	ccattcttaa	tttaactatt	tatattatcc	5100
taactactac	cgcatccta	ctactcaact	taaactccag	caccacgacc	ctactactat	5160
ctcgcacctg	aaacaagcta	acatgactaa	cacccttaat	tccatccacc	ctcctctccc	5220
taggaggcct	gcccccgcta	accggctttt	tgcccaaagt	ggccattatc	gaagaattca	5280
caaaaaacaa	tagcctcctc	atccccacca	tcatagccac	catcaccttc	cttaacctct	5340
acttctacct	acgcctaata	tactccacct	caatcacact	actccccata	tctaacaacg	5400
taaaaaataaa	atgacagttt	gaacatacaa	aacccacccc	attcctcccc	acactcatcg	5460
cccttaccac	gctactccta	cctatctccc	cttttatact	aataatctta	tagaaattta	5520
ggtaaatac	agaccaagag	ccttcaaagc	cctcagtaag	tgcaataact	taatttctgt	5580
aacagctaag	gactgcaaaa	ccccactctg	catcaactga	acgcnaatca	gccactttta	5640
traagctaag	cccttactag	accaatggga	cttaaaccca	caaacactta	gttaacagct	5700
aagcacccta	atcaactggc	ttcaatctac	ttctcccgc	gccgggaaaa	aaggcgggag	5760
aagccccggc	aggtttgaag	ctgcttcttc	gaatttgcaa	ttcaatatga	aaatcacctc	5820
ggagctggta	aaaagaggcc	taaccctgt	ctttagattt	acagtcfaat	gcttcactca	5880
gccattttac	ctcaccccca	ctgatgttcg	ccgaccgttg	actattctct	acaaaccaca	5940
aagacattgg	aacactatac	ctattattcg	gcgcagtagc	tgagtccta	ggcacagctc	6000
taagcctcct	tattcgagcc	gagctgggcc	agccaggcaa	ccttctaggt	aacgaccaca	6060
tctacaacgt	tatcgtcaca	gcccattgat	ttgtaataat	cttcttcata	gtaataacca	6120
tcataatcgg	aggctttggc	aactgactag	ttccccta	aatcggtgcc	ccgatattgg	6180
cgtttccccg	cataaacaac	ataagcttct	gactcctacc	tccctctctc	ctactcctgc	6240
tcgcatctgc	tatagtggag	gccggagcag	gaacagggtg	aacagtctac	cctcccttag	6300
cagggaacta	ctcccaccct	ggagcctccg	tagacctaac	catcttctcc	ttacacctag	6360
caggtgtctc	ctctatctta	ggggccatca	atttcatcac	aacaattatc	aatataaaac	6420
cccctgccc	aacccaatac	caaacgcccc	tcttcgtctg	atccgtccta	atcacagcag	6480
tcctacttct	cctatctctc	ccagtcctag	ctgctggcat	cactatacta	ctaacagacc	6540
gcaacctcaa	caccaccttc	ttcgaccccg	ccggaggagg	agacccatt	ctataccaac	6600
acctattctg	atttttcggg	caccctgaag	tttatattct	tatcctacca	ggcttcggaa	6660
taatctccca	tattgtaact	tactactccg	gaaaaaaaga	accatttgga	tacataggta	6720
tggtctgagc	tatgatatca	attggcttcc	tagggtttat	cgtgtgagca	caccatatat	6780
ttacagtagg	aatagacgta	gacacacgag	catatttcac	ctccgctacc	ataatcatcg	6840
ctatccccac	cggcgtcaaa	gtatttagct	gactcgccac	actccacgga	agcaatatga	6900
aatgatctgc	tgagtgctc	tgagccctag	gattcatctt	tcttttcacc	gtagggtggc	6960
tgactggcat	tgtattagca	aactcatcac	tagacatcgt	actacacgac	acgtactacg	7020
ttgtagccca	cttccactat	gtcctatcaa	taggagctgt	atttgccatc	ataggaggct	7080
tcattcactg	atttccccta	ttctcaggct	acaccctaga	ccaaaacctac	gcaaaaatcc	7140
atttcactat	catattcatc	ggcgtaaatc	taactttctt	cccacaacac	tttctcggcc	7200

tatccggaat	gccccgacgt	tactcggact	accccgatgc	atacaccaca	tgaaacatcc	7260
tatcatctgt	aggctcattc	atttctctaa	cagcagtaat	attaataatt	ttcatgattt	7320
gagaagcctt	cgcttcgaag	cgaaaagtcc	taatagtaga	agaaccctcc	ataaacctgg	7380
agtgactata	tggatgcccc	ccaccctacc	acacattcga	agaaccctga	tacataaaat	7440
ctagacaaaa	aaggaaaggaa	tgaaccccc	caaagctggt	ttcaagccaa	ccccatggcc	7500
tccatgactt	tttcaaaaag	gtattagaaa	aaccattttca	taactttgtc	aaagttaaat	7560
tataggctaa	atcctatata	tcttaatggc	acatgcagcg	caagtaggtc	tacaagacgc	7620
tacttccccct	atcatagaag	agcttatcac	ctttcatgat	cacgccccca	taatcatttt	7680
ccttatctgc	ttcctagtcc	tgtatgccct	tttccctaaca	ctcacaacaa	aactaactaa	7740
tactaacatc	tcagacgctc	aggaaataga	aaccgtctga	actatcctgc	ccgccatcat	7800
cctagtccctc	atcgccctcc	catccctacg	catcctttac	ataacagacg	aggtaacga	7860
tccctccctt	accatcaaat	caattggcca	ccaatggtac	tgaacctacg	agtacaccga	7920
ctacggcgga	ctaactcttca	actcctacat	acttccccca	ttattcctag	aaccaggcga	7980
actgcgactc	cttgacgttg	acaatcgagt	agtactcccc	attgaagccc	ccattcgtat	8040
aataattaca	tcacaagacg	tcttgcactc	atgagctgtc	cccacattag	gcttaaaaaac	8100
agatgcaatt	cccggacgtc	taaaccaaac	cactttcacc	gctaacgacg	cgggggtata	8160
ctacgggtcaa	tgctctgaaa	tctgtggagc	aaaccacagt	ttcatgcccc	tcgtcctaga	8220
attaattccc	ctaaaaatct	ttgaaatagg	gcccgatttt	accctatagc	acccccctca	8280
ccccctctag	agcccactgt	aaagctaact	tagcattaac	cttttaagtt	aaagattaag	8340
agaaccaaca	cctctttaca	gtgaaatgcc	ccaactaaat	actaccgtat	ggcccacccat	8400
aattaccccc	atactcctta	cactattcct	catcacccaa	ctaaaaatat	taaacacaaa	8460
ctaccaccta	cctccctcac	caaagcccat	aaaaataaaa	aattataaca	aaccctgaga	8520
accaaaatga	acgaaaatct	gttcgcttca	ttcattgccc	ccacaatcct	aggcctaccc	8580
gccgcagtag	tgatcattct	atttccccct	ctattgatcc	ccacctccaa	atatctctac	8640
aacaaccgac	taatcaccac	ccaacaatga	ctaatacaac	taacctcaaa	acaatgata	8700
accatacaca	acactaaagg	acgaacctga	tctcttatac	tagtatccct	aatcattttt	8760
attgccacaa	ctaacctcct	cggactcctg	cctcactcat	ttacaccaac	cacccaacta	8820
tctataaacc	tagccatggc	catcccccta	tgagcgggca	cagtgattat	aggctttcgc	8880
tctaagatta	aaaatgccct	agcccacttc	ttaccacaag	gcacacctac	accccttata	8940
cccatactag	ttattatcga	aaccatcagc	ctactcattc	aaccaatagc	cctggccgta	9000
cgcttaaccg	ctaacattac	tgcaggccac	ctactcatgc	acctaattgg	aagcgccacc	9060
ctagcaatat	caaccattaa	ccttccctct	acacttatca	tcttcacaat	tctaattcta	9120
ctgactatcc	tagaaaatcg	tgtcgcctta	atccaagcct	acgtttttcac	acttctagta	9180
agcctctacc	tgcacgacaa	cacataatga	cccaccaatc	acatgcctat	catatagtaa	9240
aaccagccc	atgaccccta	acagggggcc	tctcagccct	cctaatagacc	tcgggcctag	9300
ccatgtgatt	tcacttccac	tccataacgc	tctctatact	aggcctacta	accaacacac	9360
taaccatata	ccaatgatgg	cgcgatgtaa	cacgagaaaag	cacataccaa	ggccaccaca	9420
caccacctgt	ccaaaaaggc	cttcgatacg	ggataatcct	atttattacc	tcagaagttt	9480
ttttcttcgc	aggatttttc	tgagcctttt	accactccag	cctagccctc	accccccaat	9540
taggagggca	ctggccccc	acaggcatca	ccccgctaaa	tcccctagaa	gtcccactcc	9600
taaacacatc	cgtattactc	gcatacggag	tatcaatcac	ctgagctcac	catagtctaa	9660
tagaaaaaca	ccgaaaacaa	ataattcaag	cactgcttat	tacaatttta	ctgggtctct	9720
attttaccct	cctacaagcc	tcagagtact	tcgagtctcc	cttcaccatt	tccgacggca	9780
tctacggctc	aacatttttt	gtagccacag	gcttccacgg	acttcacgtc	attattgggt	9840
caactttcct	cactatctgc	ttcatccgcc	aactaatatt	tcactttaca	tccaaacatc	9900
actttggctt	cgaagccgcc	gcctgatact	ggcattttgt	agatgtgggt	tgactatttc	9960
tgtatgtctc	catctattga	tgagggtctt	actcttttag	tataaatagt	accgttaact	10020
tccaattaac	tagttttgac	aacattcaaa	aaagagtaat	aaacttcgcc	ttaattttta	10080
taatcaacac	cctcctagcc	ttactactaa	taattattac	attttgacta	ccacaactca	10140
acggctacat	agaaaaatcc	accccttacg	agtgcggctt	cgaccctata	tcccccgccc	10200
gcgtcccttt	ctccataaaa	ttcttcttag	tagctattac	cttcttatta	tttgatctag	10260
aaattgccc	ccttttacc	ctaccatgag	ccttacaac	aactaacctg	ccactaatag	10320
ttatgtctac	cctcttatta	atcatcatcc	tagcccttaag	tctggcctat	gagtgactac	10380
aaaaaggatt	agactgaacc	gaattgggat	atagttttaa	caaaacgaat	gatttcgact	10440
cattaaatta	tgataatcat	atttaccaaa	tgccctcat	ttacataaat	attatactag	10500



catttaccat	ctcacttcta	ggaatactag	tatatcgctc	acacctcata	tcctccctac	10560
tatgcctaga	aggaataata	ctatcgctgt	tcattatagc	tactctcata	accctcaaca	10620
cccactccct	cttagccaat	attgtgccta	ttgccatact	agtctttgcc	gcctgcgaag	10680
cagcgggtggg	cctagcccta	ctagtctcaa	tctccaaacac	atatggccta	gactacgtac	10740
ataacctaata	cctactccaa	tgctaaaact	aatcgtccca	acaattatat	tactaccact	10800
gacatgactt	tccaaaaaac	acataatttg	aatcaacaca	accaccaca	gcctaattat	10860
tagcatcatc	cctctactat	tttttaacca	aatcaacaac	aacctattta	gctgttcccc	10920
aacctttttc	tccgaccccc	taacaacccc	cctccataata	ctaactacct	gactcctacc	10980
cctcacaatc	atggcaagcc	aacgccactt	atccagtga	ccactatcac	gaaaaaaact	11040
ctacctctct	atactaactc	ccctacaaat	ctccttaatt	ataacattca	cagccacaga	11100
actaatcata	ttttatatct	tcttcgaaac	cacacttate	cccaccttgg	ctatcatcac	11160
ccgatgaggg	aaccagccag	aacgcctgaa	cgcaggcaca	tacttcctat	tctacaccc	11220
agtaggctcc	cttccccctac	tcctcgactc	aattttacact	cacaacaccc	taggctcact	11280
aaacattcta	ctactcactc	tcactgcccc	agaactatca	aactcctgag	ccaataactt	11340
aatatgacta	gcttacacaa	tagcttttat	agtaaagata	cctctttacg	gactccactt	11400
atgactccct	aaagccccatg	tcgaagcccc	catcgctggg	tcaatagtac	ttgccgcagt	11460
actcttaaaa	ctagggcggt	atgggtataat	acgcctcaca	ctcattctca	acccctgac	11520
aaaacacata	gcctaccctt	tccttgtact	atccctatga	ggcataatta	taacaagctc	11580
catctgccta	cgacaaaacag	acctaaaatc	gctcattgca	tactcttcaa	tcagccacat	11640
agccctcgta	gtaacagcca	ttctcatcca	aaacccctga	agcttcaccg	gcgcagtcac	11700
tctcataatc	gcccacgggc	ttacatcctc	attactattc	tgcttagcaa	actcaaacta	11760
cgaacgcact	cacagtcgca	tcataatcct	ctctcaagga	cttcaaactc	tactcccact	11820
aatagctttt	tgatgacttc	tagcaagcct	cgctaacctc	gccttacccc	ccactattaa	11880
cctactggga	gaactctctg	tgctagtaac	cacgttctcc	tgatcaaata	tcactctcct	11940
acttacagga	ctcaacatac	tagtcacagc	cctatactcc	ctctacatat	ttaccacaac	12000
acaatggggc	tcactcacc	accacattaa	caacataaaa	ccctcattca	cacgagaaaa	12060
cacctcatg	ttcatacacc	tatccccat	tctcctccta	tcctccaacc	ccgacatcat	12120
taccgggttt	tcctcttgta	aatatagttt	aaccaaaaca	tcagattgtg	aatctgacaa	12180
cagaggctta	cgacccctta	tttaccgaga	aagctcacia	gaactgctaa	ctcatgcccc	12240
catgtctaac	aacatggctt	tctcaacttt	taaaggataa	cagctatcca	ttggctcttag	12300
gccccaaaaa	ttttggtgca	actccaaata	aaagtaataa	ccatgcacac	tactataacc	12360
accctaacc	tgacttccct	aattcccccc	atccttacca	ccctcgtaa	ccctaacaaa	12420
aaaaactcat	acccccatta	tgtaaaatcc	attgtcgcat	ccaccttat	tatcagtctc	12480
ttccccacaa	caatattcat	gtgcctagac	caagaagtta	ttatctcgaa	ctgacactga	12540
gccacaaccc	aaacaacca	gctctcccta	agcttcaaac	tagactactt	ctccataata	12600
ttcatccctg	tagcattgtt	cgttacatgg	tcctcatag	aattctcact	gtgatataa	12660
aactcagacc	caaacattaa	tcagttcttc	aaatatctac	tcacttccct	aattaccata	12720
ctaactcttag	ttaccgctaa	caacctattc	caactgttca	tcggctgaga	gggcgtagga	12780
attatatcct	tcttgctcat	cagttgatga	tcagcccgag	cagatgcaa	cacagcagcc	12840
attcaagcaa	tcctatacaa	ccgtatcggc	gatatcggtt	tcctcctcgc	cttagcatga	12900
tttatcctac	actccaactc	atgagaccca	caacaaatag	cccttctaaa	cgctaatacca	12960
agcctcacc	cactactagg	cctcctccta	gcagcagcag	gcaaatacag	ccaattaggt	13020
ctccacccct	gactccctc	agccatagaa	ggccccaccc	cagtctcagc	cctactccac	13080
tcaagcacta	tagttgtagc	aggaatcttc	ttactcatcc	gcttccaccc	cctagcagaa	13140
aatagcccac	taatccaaac	tctaactata	tgcttaggcg	ctatcaccac	tctgttcgca	13200
gcagtctgcg	cccttacaca	aaatgacatc	aaaaaaatcg	tagccttctc	cacttcaagt	13260
caactaggac	tcataatagt	tacaatcggc	atcaaccaac	cacacctagc	attcctgcac	13320
atctgtaccc	acgccttctt	caaagccata	ctattttatgt	gctccgggtc	catcatccac	13380
aaccttaaca	atgaacaaga	tattcgaaaa	ataggaggac	tactcaaaac	catacctctc	13440
acttcaacct	ccctcaccat	tggcagccta	gcattagcag	gaataacctt	cctcacaggt	13500
ttctactcca	aagaccacat	catcgaaacc	gcaaacatat	catacacaaa	cgctgagcc	13560
ctatctatta	ctctcatcgc	tacctccctg	acaagcgctt	atagcactcg	aataattctt	13620
ctcaccctaa	caggtcaacc	tcgcttcccc	acccttacta	acattaacga	aaataacccc	13680
accctactaa	accccatata	acgcctggca	gccggaagcc	tattcgaggg	atttctcatt	13740
actaacaaca	tttcccccg	atcccccttc	caaacaacaa	tcctcctcta	cctaaaaactc	13800

```

acagccctcg ctgtcacttt cctaggactt ctaacagccc tagacctcaa ctacctaacc 13860
aacaaactta aaataaaatc cccactatgc acattttatt tctccaacat actcggattc 13920
taccctagca tcacacaccg cacaatcccc tatctaggcc ttcttacgag ccaaaacctg 13980
cccctactcc tcctagacct aacctgacta gaaaagctat tacctaaaac aatttcacag 14040
caccaaactc ccacctccat catcacctca acccaaaaag gcataattaa actttacttc 14100
ctctctttct tcttcccact catcctaacc ctactcctaa tcacataacc tattcccccg 14160
agcaatctca attacaatat atacaccaac aaacaatgtt caaccagtaa ctactactaa 14220
tcaacgccc aatacatata aagccccgc accaatagga tcctcccgaa tcaacctga 14280
ccccctctct tcataaatta tttagcttcc tacactatta aagtttacca caaccaccac 14340
cccatcatat tctttcacc acagcaccaa tcctacctcc atcgctaacc cactaaaaac 14400
actcaccaag acctcaaccc ctgaccccc tgctcagga tactcctcaa tagccatcgc 14460
tgtagtatat ccaaagacaa ccatcattcc ccctaaataa attaaaaaaa ctattaaacc 14520
catataacct cccccaaaat tcagaataat aacacaccgc accacaccgc taacaatcaa 14580
tactaaaccc ccataaatag gagaaggctt agaagaaaac ccacaaaacc ccattactaa 14640
accacactc aacagaaaca aagcatatat cattattctc gcacggacta caaccacgac 14700
caatgatatg aaaaaccatc gttgtatttc aactacaaga acaccaatga ccccaatacg 14760
caaaattaac cccctaataa aattaattaa cactcattc atcgacctcc ccaccccatc 14820
caacatctcc gcatgatgaa acttcggctc actccttggc gcctgcctga tcctccaaat 14880
caccacagga ctattcctag ccattgcacta ctaccagac gcctcaaccg ccttttcac 14940
aatcgccac atcactcgag acgtaaatta tggctgaatc atccgctacc ttcacgcaa 15000
tggcgctca atattcttta tctgcctct cctacacatc gggcgaggcc tatattacgg 15060
atcatttctc tactcagaaa cctgaaacat cggcattatc ctctgcttg caactatagc 15120
aacagccttc ataggctatg tcctcccgat aggccaaata tcattctgag gggccacagt 15180
aattacaaac ttactatccg ccatcccata cattgggaca gacctagttc aatgaatctg 15240
aggaggctac tcagtagaca gtcccacct caccagattc ttacctttc acttcatctt 15300
gcccttcatt attgcagccc tagcaacact ccacctccta ttcttgacg aaacgggatc 15360
aaacaacccc ctaggaatca cctcccattc cgataaaatc acctccacc ctactacac 15420
aatcaaagac gccctcggtc tacttctctt ccttctctcc ttaatgacat taacactatt 15480
ctcaccagac ctccctaggcg acccagacaa ttatacccta gccaacccct taaacacccc 15540
tccccacatc aagcccgaa gatatttctt attcgctac acaattctcc gatccgtccc 15600
taacaaacta ggaggcgctc ttgccctatt actatccatc ctcatcctag caataatccc 15660
catcctccat atatccaaac aacaaagcat aatatttctc ccactaagcc aatcacttta 15720
ttgactccta gccgcagacc tcctcattct aacctgaatc ggaggacaac cagtaagcta 15780
cccttttacc atcattggac aagtagcatc cgtactatac ttcacaacaa tcctaactct 15840
aataccaact atctccctaa ttgaaaacaa aatactcaa tgggcctgtc cttgtagtat 15900
aaactaatc accagtcttg taaaccggag atgaaaacct tttccaagg acaaatcaga 15960
gaaaaagtct ttaactccac cattagcacc caaagctaag attctaattt aaactattct 16020
ctgttctttc atggggaagc agatttgggt accaccaag tattgactca ccatcaaca 16080
accgctatgt atttctgata ttactgccag ccaccatgaa tattgtacgg taccataaat 16140
acttgaccac ctgtagtaca taaaaaccca atccacatca aaacccctc ccatgctta 16200
caagcaagta cagcaatcaa ccctcaacta tcacacatca actgcaactc caaagccacc 16260
cctcaccac taggatacca acaaacctac ccacccttaa cagtacatag tacataaagc 16320
catttaccgt acatagcaca ttacagtcaa atcccttctc gtcccatgg atgaccccc 16380
tcagataggg gtcccttgac caccatctc cgtgaaatca atatcccgca caagagtgtc 16440
actctcctcg ctccggggcc ataacacttg ggggtagcta aagtgaactg tatccgacat 16500
ctggttctca cttcagggc ataaagccta aatagccac acgttcccct taaataagac 16560
atcacgatg

```

&lt;210&gt; 3

&lt;211&gt; 6691

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(6691)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 3

ataggcattc	caactcatcc	gcttgctgac	gacctccacg	tgatttcaac	aatgatttca	60
aatatattac	tttttaagtc	agtgtgactt	aagtatgaaa	ttgcctctcc	ctaaagctcc	120
cctaaggcct	aaacagtcgt	cattaccata	gctgtgacag	ggagactggt	gaatttataa	180
tctattggcc	attcacagca	tagcgtataa	accctagctc	atgatttctt	tgcaatagaa	240
gtgtactttt	tcatcacatt	cccttcacaa	cttactcacc	agacagactt	tgagctctcc	300
tcctggccta	gcctggatcg	tttgaaatgg	tcattccatcc	tttggcccca	atacctaaac	360
taaggcttat	gaacaataag	atgattttac	ttcagtggga	cttttttggt	taatatatta	420
gatttgacct	tcagcaagg	caaagggagt	ccgaactagt	ctcaggcttc	aacatcgaat	480
acgccgcagg	ncccttcgcc	ctattcttca	tagccgaata	cacaaacatt	attataataa	540
acaccctcac	cactacaatc	ttcctaggaa	caacatataa	cgcactctcc	cctgaactct	600
acacaacata	ttttgtcacc	aagaccctac	ttctgacctc	cctgttctta	tgaattcgaa	660
cagcatcccc	ccgattccgc	tacgaccaac	tcataacctt	cctatgaaaa	aacttcctac	720
cactcacctt	agcattactt	atatgatatg	tctccatacc	cattacaatc	tccagcattc	780
cccccaaac	ctaagaaata	tgtctgataa	aagagttact	ttgatagagt	aaataatagg	840
agtttaaatc	cccttatttc	taggactatg	agaatcgaac	ccatccctga	gaatccaaaa	900
ttctcggtgc	caactatcac	accccatctt	aaagtaagg	cagctaaata	agctatcggg	960
cccatacccc	gaaaaatgtg	gttatatcct	tcccgtaacta	attaatcccc	tggcccaacc	1020
cgtcatttac	tctaccatct	ttgcaggcac	actcatcaca	gcgctaagct	cgcactgatt	1080
ttttacctga	gtaggcctag	aaataaacat	gctagrtttt	attccagttc	taacccaaaa	1140
aataaacctt	cgttccacag	aagntgccat	caagtatttc	ctcacgcaag	caaccgcac	1200
cataatcctt	ctaatagcta	tcctnttcaa	caataatact	tcgggacaat	gaaccatwac	1260
caataccacc	aatcaatact	catcattaat	aatcataatg	gctatagcaa	taaaactagg	1320
aatagccccc	tttcaacttc	gagtcacaga	ggttacccaa	ggcacccttc	tgacatccgg	1380
cctgctcctt	ctcacatgac	aaaaactagc	ccccatctca	atcatatacc	aaatttctcc	1440
ctcattaaac	gtaagccttc	tcctcactct	ttcaatctta	tccatcatgg	caggcagttg	1500
agggtggatta	aaccaaaacc	aactacgcaa	aatcttagca	tactcctcaa	ttaccacat	1560
aggatgaata	acagcagttc	taccgtacaa	ccctaacata	accattctta	atttaactat	1620
ttatattatc	ctaactacta	ccgcattcct	actactcaac	ttaaactcca	gcaccacaac	1680
cctactacta	tctcgcacct	gaaacaagct	aacatgacta	acacccttaa	ttccatccac	1740
cctcctctcc	ctaggaggcc	tgcccccgct	aaccggcttt	ttgcccgaat	gggccattat	1800
cgaagaattc	acaaaaaaca	atagcctcat	catccccacc	atcatagcca	tcataccctt	1860
ccttaacctc	tacttctacc	tgcgccctaat	ctactccacc	tcaatcacac	tactccctat	1920
atctaacaac	gtaaaaataa	aatgacagtt	tgaacacaca	aaaccacccc	cattcctccc	1980
cacactcatc	gcccttacca	cactgctcct	acctatctcc	ccttttatgc	taataatctt	2040
atagaaattt	aggttaaata	cagaccaaga	gccttcaaag	ccctcagtaa	gttgcaatac	2100
ttaatttctg	naacagctaa	ggactgcaaa	acccactctt	gcatacaactg	nangcaaatc	2160
agccacttta	attaagctaa	gcccttacta	gaccaatggg	acttaaacc	acaaacactt	2220
agttaacagc	taagcaccct	aatcaactgg	cttcaatcta	cttctcccgc	cgccgggaaa	2280
aaaggcggga	gaagccccgg	cagggttgaa	gctgcttctt	cgaatttgca	attcaatatg	2340
aaaatcacct	canagctggt	aaaaagaggc	ttaacccttg	tcttttagatt	tacagtccaa	2400
tgcttcactc	agccatttta	cctcaccccc	actgatgttc	gccgaccgtt	gactattctc	2460
tacaaaccac	aaagacattg	gaacactata	cctattatct	ggcgcatgag	ctggagtctt	2520
aggcacagct	ctaagcctcc	ttattcgagc	cgaactgggc	cagccaggca	accttctagg	2580
taacgaccac	atctacaacg	ttatcgtcac	agcccatgca	tttgtaataa	tcttcttcat	2640
agtaataccc	atcataatcg	gaggytttgg	caastgacta	gttcccctaa	taatcggtgc	2700
ccccgatatg	gsgtttcccc	gcataaacia	cataagcttn	tgactcttac	ccccctctct	2760
cntactcctg	ttngcatctg	ctatagtggg	ggccggcgca	ggaacagggt	gaacagnta	2820
ccctcccttg	gcagggaact	actcccaccc	tggagcctcc	gtagacctaa	ccatcttctc	2880
cttacacctt	gcagggtatct	cctctatctt	agggaccatc	aatttcatca	caacaattat	2940
taatataaaa	ccccctgcca	taaccaata	ccaaacgccc	cttttcgtct	gatccgtcct	3000
aatcacagca	gtcttacttc	tcctatctct	cccagtccta	gccgctggca	tcactatact	3060

actaacagac	cgtaacctca	acaccacctt	cttcgaccca	gccggaggag	gagaccccat	3120
tctataccaa	cacctattct	gatttttcgg	tcaccctgaa	gtttatattc	tcacccctacc	3180
aggcttcgga	ataatctccc	atattgtaac	ttactactcc	ggaaaaaaag	aaccatttgg	3240
atacataggt	atggctctgag	ctatgatatc	aattggcttc	ctagggttta	tcgtgtgagc	3300
acaccatata	tttacagtag	gaatagacgt	agacacacga	gcatatttca	cctccgctac	3360
cataatcatc	gctatcccca	ccggcgctca	agtatttagc	tgactcgcca	cactccacgg	3420
aagcaatâtg	aaatgatctg	ctgcagtgtc	ctgagcccta	ggatttattt	ttcttttcac	3480
cgtagggtggc	ctgactggca	ttgtatttagc	aaactcatca	ctagacatcg	tactacacga	3540
cacgtactac	gttgtagccc	acttccacta	tgtcctatca	ataggagctg	tatttgccat	3600
cataggaggc	ttcattcact	gatttcccct	attctcaggc	tacaccctag	accaaacctta	3660
cgccaaaatc	catttcgcta	tcataattcat	cggcgtaaat	ctaactttct	tcccacaaca	3720
ctttctcggc	ctatccggaa	tgccccgacg	ttactcggac	tatcccgatg	catacaccac	3780
atgaaatatc	ctatcatctg	taggctcatt	catttctcta	acagcagtaa	tattaataat	3840
tttcataatt	tgagaagcct	tcgcttcgaa	gcgaaaagtc	ctaatagtag	aagaaccttc	3900
cataaacctg	gagtgtactat	atggatgccc	cccacctac	cacacattcg	aagaacctcg	3960
atacataaaa	tctagacaaa	aaaggaagga	atcgaacccc	ccaaagctgg	tttcaagcca	4020
accccatggc	ctccatgact	ttttcaaaaa	gatattagaa	aaaccatttc	ataactttgt	4080
caaagttaaa	ttataggcta	aatcctatat	atcttaatgg	cacatgcagc	gcaagtaggt	4140
ctacaagacg	ctacttcccc	tatcatagaa	gagcttatca	tctttcatga	tcacgccttc	4200
ataatcattt	tccttatctg	cttcctagtc	ctgtacgccc	ttttcctaac	actcacaaca	4260
aaactaacta	atactaacta	ctcagacgct	caggaaatag	aaaccgtctg	aactatcctg	4320
cccgccatca	tcctagtcct	tatcgccctc	ccatnccctac	gcacccctta	cataacagac	4380
gagggtcaacg	atccctcctt	taccatcaaa	tcaattggcc	atcaatggta	ctgaacctac	4440
gaatacaccg	actacggcgg	actaatcttc	aactctaca	tacttcccc	attatttcta	4500
gaaccaggcg	acctgcgact	cccttgacgtt	gacaactcag	tagtactccc	gggtgaagcc	4560
cccattcgtâ	taataattac	atcacaagac	gtcttacact	catgagctgt	ccccacatta	4620
ggcttaaaaa	cagatgcaat	tcccgacgt	ctaaaccaa	ccactttcac	tgctacacga	4680
ccaggggtat	actacggcca	atgctctgaa	atctgtggag	caaaccagtt	ttatgcccat	4740
cgtcctagaa	ttaattcccc	taaaaatctt	tgaaataggg	cccgtattta	ccctatagca	4800
ccccctctac	ccccctctaga	gcccactgta	aagctaactt	agcattaacc	ttttaagtta	4860
aagattaaga	gaaccaacac	ctcttttacag	tgaaatgccc	caactaaata	ctaccgtatg	4920
accaccata	attaccccca	tactccttac	actattcctc	atcacccaac	taaaaatatt	4980
aaatacaaat	taccacctac	ctccctcacc	aaagcccata	aaaataaaaa	actataacaa	5040
accctgagaa	ccaaaatgaa	cgaaaatctg	ttcacttcat	tcattgcccc	cacaatccta	5100
ggcctaccgg	ccgcagtact	gatcatttcta	ttycccccctc	tattgatccc	cacctccaaa	5160
tatctcatca	acaaccgact	aattaccacc	caacaatgac	taatccaact	aacctcaaaa	5220
caaatgatag	ccatacacaa	cactaaggga	cgaacctgat	ctcttatact	agtatcctta	5280
atcattttta	ttgccacaac	taacctcctc	ggactcctgc	ctcactcatt	tacaccaacc	5340
acccaactat	ctataaacct	agccatggcc	atcccccttat	gagcgggcgc	agtgattata	5400
ggctttcgtc	ctaagattaa	aaatgcccta	gcccacttct	taccacaagg	cacacctaca	5460
ccccctatcc	ctatactagt	tattatcgaa	accatcagcc	tactcattca	accaatagcc	5520
ctggccgtan	gnctaaccgc	taacattact	gcaggccacc	tactcatgna	cctaattgga	5580
agcgccacac	tagcaatatc	aantattaac	cttccctcta	cacttatcat	tttcacaatt	5640
ctaattctac	tgactatcct	agaaatcgct	gtcgccttaa	tccaagccta	cgtttttaca	5700
cttttagtaa	gcctctacct	gcacgacaac	acataatgac	ccaccaatca	catgcctatc	5760
atatagtaaa	acccagccca	tggccccctaa	caggggccct	ctcancctc	ctaatagcct	5820
ccggcctagc	catgtgattt	cacttccact	ccacaaccct	ccttatacta	ggcctactaa	5880
ccaacacact	aaccatatac	caatgatggc	gcgatgtaac	acgagaaagc	acataccaag	5940
gccaccacac	accacctgtc	cagaaaggcc	ttcgatacgg	gataatccta	tttattacct	6000
cagaagtttt	tttcttcgca	ggatttttct	gagcctttta	ccactccagc	ctagccccna	6060
ccccccaact	agggggacac	tggcccccaa	caggcatcac	cccgtctaat	cccctagaag	6120
tcccactcct	aaacacatcc	gtattactcg	catcaggggt	atcaatcacc	tgagctcacc	6180
atagtcta	agaaaacaa	cgaaaacaaa	taattcaagc	actgcttatt	acaattttac	6240
tgggtctcta	ttttaccctc	ctacaagcct	cagngtactt	cgagggttaa	atattagata	6300
tttccctga	tacagggtc	aatctttttc	tttttaagc	aatatttctc	aaagtacttt	6360

tcacagaact taagtttcat taagcacttc actaaaagna aaagtctgtg atctaataaa	6420
tttggaatat attgagaatt agagccccct cttagatatg tactgtagct actcagcttg	6480
ttacagatgg aagtaaacat tgtaatatcc acccagcttt tgagtggatg tctattaaca	6540
tcaccccaat gagtattcca tggaatgcac ttgcaaaaaa cctattattc aagaaaaatt	6600
ctggagcatg gaaagctatt aatggataaa cccattcaca aaatcacacc aaatatctaa	6660
aatcatgttt aaaatctcct agaaatgggt t	6691

&lt;210&gt; 4

&lt;211&gt; 1263

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 4

agagagagggc atgtgaattg ggaatttggg aaaaattttt tggggggaag gaaagaaata	60
gaggtcaaga ggtagaatag aagttgatga agaaaagaaa aaaagaagggt aatgaagggg	120
gtgctggatg ttccaacac aaagaaatga taaatgtttg ggaggatgga tattctaatt	180
agcctaatta gcctgattag ccctcgccag agttcactgt aaagctaacc cagcattaac	240
cttttaagtt aaagactaag agaatcatta tctctttaca gtgaaatgcc acagctaaat	300
accactgtat gacctgctat catcacccca atactcctca cgttatttct catcacccaa	360
ctaaaaatac taaacacaca ctgccatctg cccacctcac caaaatttat taaaataaaa	420
aactacagta agccctgaga accaaaatga acgaaaattt attcgcttca ttcattaccc	480
ctacagtact aggcctaccc gccacagtac caatcatcct atttcccccc ttactgggcc	540
caacctccaa atacctcatc aacaaccgac taatcaccac tcaacaatga ctacttcaac	600
tcaccttaaa acaataata acgatacata acattaaggg acgaacctgg tcccttatac	660
taatttccct gattattttt attgccacaa ctaatctcct cggactcttg cccactcat	720
ttacaccaat cactatacat gtgtctattg aaacgtcact atgtgtgccc catgaatatg	780
tacatattat tatgtgatgt acatgattat gtacacatta tgtgccaatg aaataaataa	840
aattaaaagg taattcccaa ataaataaaa acagaaaatt ttaaaaagag agagagagag	900
atgagatggg gctttgggtg ggaacctggg tatggaaatc tgctgctgcc ctctttccct	960
taggaatgca agactttaat tatctccata ccaactgctt tgagatcacg ctggaactga	1020
gttgcgacaa gtttcccccc gaagaggagt tacagcggga gtggctgggt aatcggggag	1080
ccctaattcca gttcctggaa caggtaaaat ctcatttgtt aactctcttg ttgcaaaaaa	1140
gagaagtgtc tctcaatggc aggcaagtgt ttcactgggt cagtctgact ttgctctga	1200
ctctgtccaa tatatctggg cgcttcttct ccattgagat tttattcatt tccaccttta	1260
ttt	1263

&lt;210&gt; 5

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Primer complementary to mitochondrial DNA  
sequences in the CO1 encoding region of SEQ ID  
NO:2

&lt;400&gt; 5

ccttacacct agcaggta

18

&lt;210&gt; 6

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Primer complementary to mitochondrial DNA  
sequences in the CO1 encoding region of SEQ ID  
NO:2

<400> 6  
acgccgatga atatgatagc 20

<210> 7  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer

<400> 7  
catgcagcgc aagtaggtct acaagac 27

<210> 8  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer

<400> 8  
tggttatgtaa aggatgcgta gggatgg 27

<210> 9  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer

<400> 9  
tatgagggcg tgatcatgaa ag 22

<210> 10  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer

<400> 10  
ggcacggctg tccaagg 17

<210> 11  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer

<400> 11  
cccggcctgg tacactg 17

<210> 12  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer

<400> 12  
gcggacatgg aggacgtg 18

<210> 13  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer

<400> 13  
cgatgccgat gacctgcaga ag 22

<210> 14  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 14  
gtccaatgct tcactcagcc a 21

<210> 15  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 15  
ttacagtcca atgcttcact c 21

<210> 16  
<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 16

tatgcgggga aacgcat

18

<210> 17

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 17

tgtttatgcg gggaaacgc

19

<210> 18

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 18

ggcaactgac tagttcccct a

21

<210> 19

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 19

aatcgaggc tttggcaact g

21

<210> 20

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA



<400> 20  
gtttggtatt gggttatggc a 21

<210> 21  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 21  
tggatttggg ttatggcagg g 21

<210> 22  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 22  
ggccatcaat ttcatacaaa 20

<210> 23  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 23  
tagcaggtgt ctctctatc tt 22

<210> 24  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 24  
atacctatgt atccaaatgg ttctt 25

<210> 25  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 25  
ccatacctat gtatccaaat ggttc 25

<210> 26  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 26  
ggaataatct cccatattgt aactt 25

<210> 27  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 27  
caggccacct acggtgaa 18

<210> 28  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 28  
gtcaggccac ctacggtg 18

<210> 29  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 29  
agtgctctga gccctaggat 20

<210> 30  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 30  
atgatctgct gcagtgctct 20

<210> 31  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 31  
attccggata ggccgaga 18

<210> 32  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 32  
tcggcgtaaa tctaactttc tt 22

<210> 33  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 33  
ggggttcgat tccttcctt 19

<210> 34  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 34

ttggcttgaa accagcttt

19

<210> 35

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 35

ttgtcaaagt taaattatag gcta

24

<210> 36

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 36

gtcaaagtta aattataggc taaa

24

<210> 37

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 37

acctcgtctg ttatgtaaag gat

23

<210> 38

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 38

tgacctcgtc tgttatgtaa agg

23

<210> 39

<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 39  
cgccatcatc ctagtcctca 20

<210> 40  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 40  
atgagtgcaa gacgtcttgt gat 23

<210> 41  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 41  
aatcgagtag tactcccgat tga 23

<210> 42  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 42  
gagtagtact cccgattgaa gcc 23

<210> 43  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 43  
gttagcttta cagtgggctc taga 24

<210> 44  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 44  
agttagcttt acagtgggct ctag 24

<210> 45  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 45  
acttctagta agcctctacc tgca 24

<210> 46  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 46  
ttctagtaag cctctacctg cacg 24

<210> 47  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 47  
aggtaataaa taggattatc ccgta 25

<210> 48  
<211> 25  
<212> DNA

<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 48  
aataaatagg attatcccgat atcga 25

<210> 49  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 49  
ccacacacca cctgtccaa 19

<210> 50  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 50  
accacacacc acctgtcca 19

<210> 51  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 51  
aagggagact cgaagtactc tga 23

<210> 52  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 52

tggtgtctcta ttttaccctc cta 23

<210> 53  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 53  
tattacaatt ttactgggtc tct 23

<210> 54  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to cytochrome  
c oxidase encoding mitochondrial DNA

<400> 54  
actagttaat tggaagttaa cggta 25

<210> 55  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 55  
gaggttcaat tcctcttctt aac 23

<210> 56  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 56  
ttgagtaaac ggctaggcta ga 22

<210> 57  
<211> 20  
<212> DNA  
<213> Artificial Sequence



<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 57  
ctggtcaacc tcaacctagg 20

<210> 58  
<211> 16  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 58  
ggcctgcggc gtattc 16

<210> 59  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 59  
ccgaagggga gtccgaa 17

<210> 60  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 60  
atcaaagtaa ctcttttatc agaca 25

<210> 61  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 61  
ttggttatac cttccccgta 20

<210> 62  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 62  
ctattcctag ttttattgct atagct 26

<210> 63  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 63  
acaatatatact ctccggacaa tg 22

<210> 64  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 64  
gaatgggttat gttagggttg tac 23

<210> 65  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 65  
agctacgcaa aatcttagca ta 22

<210> 66  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH

## dehydrogenase encoding mitochondrial DNA

<400> 66  
agggtgatgg tggctatga 19

<210> 67  
<211> 15  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 67  
cccgctaacc ggctt 15

<210> 68  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 68  
agggttttga aggtctt 18

<210> 69  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 69  
tttcgactca ttaaattatg ataata 25

<210> 70  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 70  
catgtcagtg gtagtaatat aattg 25

<210> 71  
<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 71

cacatatggc ctgactacg ta

22

<210> 72

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 72

ataattaagg agatttgtag gga

23

<210> 73

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 73

ccaacgccac ttatccagt

19

<210> 74

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 74

aagctattgt gtaagctagt catat

25

<210> 75

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 75  
ctcactaaac attctactac tcact 25

<210> 76  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 76  
gtggctgatt gaagagtatg c 21

<210> 77  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 77  
cctatgaggc ataattataa caag 24

<210> 78  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 78  
acgtggttac tagcacagag ag 22

<210> 79  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 79  
tgacttctag caagcctcg 19

<210> 80  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 80  
acaatctgat gttttggtta aac 23

<210> 81  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 81  
ctcatcctaa ccctactcct aa 22

<210> 82  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 82  
gcgatggcta ttgaggagt 19

<210> 83  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 83  
gctaacccca ctaaaacact 20

<210> 84  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 84  
ttcatatcat tggtcgtggt 20

<210> 85  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 85  
tagtaccggtt aacttccaat taact 25

<210> 86  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 86  
tcataattta atgagtcgaa atca 24

<210> 87  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 87  
cagctatcca ttggtcttag g 21

<210> 88  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH dehydrogenase encoding mitochondrial DNA

<400> 88  
tatttgaaga actgattaat gtttg 25

<210> 89  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 89  
agcattgttc gttacatggt c 21

<210> 90  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 90  
ggcttggatt agcgtttaga a 21

<210> 91  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 91  
tcacccctcgc cttagcatga 20

<210> 92  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 92  
ttttgatgtc attttgtgta agg 23

<210> 93  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 93  
caaactctaa cactatgctt agg 23

<210> 94



<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 94  
atgtttgcgg ttctgatgat 20

<210> 95  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 95  
cattggcagc ctagcattag 20

<210> 96  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 96  
gattttattt taagtttggt ggtag 26

<210> 97  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 97  
aaactcacag ccctcgctg 19

<210> 98  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to NADH  
dehydrogenase encoding mitochondrial DNA

<400> 98  
tatgtgatta ggagtagggt tagg 24

<210> 99  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to ATP  
synthetase subunits 8/6 encoding mitochondrial DNA

<400> 99  
gcccactgta aagctaactt agc 23

<210> 100  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to ATP  
synthetase subunits 8/6 encoding mitochondrial DNA

<400> 103  
tagtcggttg ttgatgagat at 22

<210> 101  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to ATP  
synthetase subunits 8/6 encoding mitochondrial DNA

<400> 101  
agtcggttgt tgatgag 17

<210> 102  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to ATP  
synthetase subunits 8/6 encoding mitochondrial DNA

<400> 102  
ggcctaccgc ccgcagtact gat 23

<210> 103  
<211> 23  
<212> DNA

<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to ATP  
synthetase subunits 8/6 encoding mitochondrial DNA

<400> 103  
tgtaggtgtg ccttggtgga aga 23

<210> 104  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to ATP  
synthetase subunits 8/6 encoding mitochondrial DNA

<400> 104  
attataggct ttcgctctaa ga 22

<210> 105  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to ATP  
synthetase subunits 8/6 encoding mitochondrial DNA

<400> 105  
ccaaccaccc aactatctat a 21

<210> 106  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to ATP  
synthetase subunits 8/6 encoding mitochondrial DNA

<400> 106  
atatgatagg catgtgattg gt 22

<210> 107  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to ATP  
synthetase subunits 8/6 encoding mitochondrial DNA

<400> 107

tagcattaac cttttaagtt aaaga 25

<210> 108  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer corresponding to ATP  
synthetase subunits 8/6 encoding mitochondrial DNA

<400> 108  
tcgttcattt tggttctca 19

<210> 109  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer for PCR amplification

<400> 109  
tccactccac aaccctcctt 20

<210> 110  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer for PCR amplification

<400> 110  
gggccagtgt cccctag 17

<210> 111  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer for PCR amplification

<400> 111  
tagcattaac cttttaagtt aaaga 25

<210> 112  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer for PCR amplification

<400> 112  
tcgttcattt tggttctca

